

# Tree Representations of Non-symmetric Group-Valued Proximities<sup>1</sup>

Charles Semple and Mike Steel

*Biomathematics Research Centre, Department of Mathematics and Statistics,  
University of Canterbury, Private Bag 4800, Christchurch, New Zealand*

E-mail: [c.semple@math.canterbury.ac.nz](mailto:c.semple@math.canterbury.ac.nz), [m.steel@math.canterbury.ac.nz](mailto:m.steel@math.canterbury.ac.nz)

Received February 10, 1999; accepted March 9, 1999

Let  $X$  be a finite set and let  $d$  be a function from  $X \times X$  into an arbitrary group  $\mathcal{G}$ . An example of such a function arises by taking a tree  $T$  whose vertices include  $X$ , assigning two elements of  $\mathcal{G}$  to each edge of  $T$  (one for each orientation of the edge), and setting  $d(i, j)$  equal to the product of the elements along the directed path from  $i$  to  $j$ . We characterize conditions when an arbitrary function  $d$  can be represented in this way, and show how such a representation may be explicitly constructed. We also describe the extent to which the underlying tree and the edge weightings are unique in such a representation. These results generalize a recent theorem involving undirected edge assignments by an Abelian group. The non-Abelian bi-directed case is of particular relevance to phylogeny reconstruction in molecular biology. © 1999 Academic Press

*Key Words:* trees; groups; four-point condition; symbolic ultrametric.

## 1. INTRODUCTION

A classical problem in classification is the following: when can an arbitrary metric on a finite set be realized by embedding the points of the metric space in a positively edge-weighted tree with its associated minimum path-length metric? More precisely, given a metric  $d: X \times X \rightarrow \mathbb{R}^{\geq 0}$ , when does there exist a tree  $T = (V, E)$  with  $X \subseteq V$  and a weighting  $w: E \rightarrow \mathbb{R}^{\geq 0}$  such that  $d(i, j)$  is the sum of the weights of the edges on the path connecting vertices  $i$  and  $j$ ? Furthermore, if  $d$  has such a representation, what can one say concerning the possible choices of  $T$  and  $w$ ?

Both questions have well-known solutions which date back 30 years (see [7, 14, 16]). Specifically, a tree representation exists for all of  $X$  precisely if

<sup>1</sup> This work was supported by the New Zealand Marsden Fund (UOC-MIS-003).



it exists for every subset of  $X$  of size at most 4, and this, in turn, is equivalent to an appropriate “four point condition” involving (in)equalities on sums of pairs of  $d(i, j)$  values. Furthermore, when they exist, the pair  $(T, w)$  that accommodates a representation of  $d$  is uniquely determined, provided  $T$  has no vertices in  $V - X$  of degree less than 3 and  $w$  is strictly positive. Note that the last two provisos can always be imposed.

These classical results, which have become a central tool in classification (particularly in evolutionary biology) have been subsequently generalized in several directions. Hakimi and Patrinos [11] considered two extensions: first, allow for edge weightings over  $\mathbb{R}$  (rather than just  $\mathbb{R}^{\geq 0}$ ); and, second, to consider trees in which each edge is assigned two real numbers (one for each orientation of the edge), with  $d(i, j)$  now being defined as the sum of the weights on the *directed* path from  $i$  to  $j$ . This second extension allows, but does not necessarily imply, non-symmetry in the function  $d$ .

A second line of generalization was adopted by Bandelt and Steel [4] to allow edge weightings to take values in a suitably structured Abelian semigroup. One spin-off of this approach was to provide a tree representation for distance hereditary graphs.

A third line of generalization was provided by Böcker and Dress [5] who developed a purely combinatorial statement (i.e., involving no algebraic structure) which implied the result of [4]; though, as pointed out in [5], the two results are actually equivalent. The main theorem from [5] will be a central tool here.

This paper represents a continuation of this story. We generalize the approach of [4] by allowing the edge weightings, and hence  $d$ , to take values in an arbitrary group, and we follow the approach of Hakimi and Patrinos of allowing each edge to have two weightings, according to its two orientations. This two-step generalization leads to only a slight complication in the statement of the main existence theorem.

A key motivation for considering these generalizations comes from the field of molecular biology, and, in particular, the problem of reconstructing evolutionary trees from aligned genetic sequences. If one assumes that these sequences evolve according to standard Markov models, then to each edge of the underlying tree is associated two transition matrices (depending on the direction along the edge that the process is run). The ordered product of these transition matrices along the path from species  $i$  to species  $j$  is then the net transition matrix for the pair  $(i, j)$  which can be estimated from genetic data (see [2, 8, 15]). Thus, if we have  $r$ -state sequences (for instance  $r = 4$  for DNA sequences), we are precisely in the setting of assigning elements of the non-Abelian group  $\mathcal{G}$  of  $r \times r$  non-singular real matrices to each orientation of the edges and taking (directed) products. The results below describe conditions under which the associated tree (and the edge-weightings) can be reconstructed (thereby generalizing

the results of [2]). Moreover, these results describe conditions under which such a representation exists over  $\mathcal{G}$  (of course, for this particular problem we require more—namely, representation over the semigroup of transition matrices, however, representation over  $\mathcal{G}$  is certainly a necessary condition). We return to this particular setting in the last section.

The structure of the paper is as follows. We begin Section 2 by setting up some terminology and establishing a basic property of tree representations. Several mappings are defined and some important relationships between these mappings are determined. In Section 3, we state the two main (existence and uniqueness) results, Theorems 3.1 and 3.2, and provide proofs. We also derive, as a corollary, the main theorem from [4]. Section 4 makes some concluding remarks.

## 2. PRELIMINARIES

Throughout this paper,  $X$  will denote a finite set, and  $\mathcal{G}$  will denote an arbitrary group with identity element  $1_{\mathcal{G}}$ . We multiply elements of  $\mathcal{G}$  from left to right.

DEFINITIONS. • Let  $T$  be a tree with vertex set  $V$  and edge set  $E \subseteq \{\{x, y\} : x, y \in V; x \neq y\}$ . A vertex  $v \in V$  is *interior* if  $\deg_T(v) > 1$ , otherwise  $v$  is a *leaf*. An edge  $e = \{u, v\} \in E$  is *interior* if both  $u$  and  $v$  are interior vertices, otherwise we say  $e$  is *exterior*.

• Suppose we have a map  $\phi: X \rightarrow V$  with the property that, for all  $v \in V$ ,

$$\deg_T(v) \leq 2 \Rightarrow v \in \phi(X).$$

The pair  $(T; \phi)$  is called an  $X$ -tree, and we sometimes write this as the ordered triple  $(V, E; \phi)$ . If  $\phi$  is a bijection from  $X$  into the set  $V_1$  of degree-one vertices of  $T$ , then  $(V, E; \phi)$  is a *phylogenetic  $X$ -tree*. In this case, we can view  $X$  as a subset of  $V_1$  and so we frequently just denote a phylogenetic  $X$ -tree by just  $T$  or  $(V, E)$ , since  $\phi$  is implicitly determined. An example of a phylogenetic  $X$ -tree for  $X = \{i, j, k, x\}$  is shown in Fig. 1. Two  $X$ -trees  $(V, E; \phi)$  and  $(V', E'; \phi')$  are *isomorphic* if there exists a bijection  $\alpha: V \rightarrow V'$  which induces a bijection between  $E$  and  $E'$  and which satisfies  $\phi' = \alpha \circ \phi$ , in which case  $\alpha$  is unique. We denote isomorphism by the symbol  $\equiv$ .

• For a tree  $(V, E)$ , let  $\vec{E} := \{(u, v) : \{u, v\} \in E\}$ . We can regard  $\vec{E}$  as the set of pairs in which each member consists of an element of  $E$  and an orientation of it. Each element of  $\vec{E}$  is called an *arc*. Let  $w$  be a function from  $\vec{E}$  into the group  $\mathcal{G}$ . We refer to  $w((u, v))$  as the *weight* of  $\text{arc}(u, v)$




FIG. 1. A phylogenetic  $X$ -tree for  $X = \{i, j, k, x\}$ .

and, for simplicity, we write  $w((u, v))$  as  $w(u, v)$ . Following [2], the *return-trip weights* of an edge  $\{u, v\}$  are the elements  $w(u, v)w(v, u)$  and  $w(v, u)w(u, v)$  of  $\mathcal{G}$ . We say that an edge  $e$  is *properly weighted* if  $1_{\mathcal{G}}$  is not a return trip weight for  $e$  (or, equivalently, if the return trip weights for  $e$  are not both equal to  $1_{\mathcal{G}}$ ).

- Given an  $X$ -tree  $(T; \phi)$  and vertices  $v_1, v_2 \in V$ , define  $d_{(T; \phi; w)}: V \times V \rightarrow \mathcal{G}$  by setting  $D_{(T; \phi; w)}(v_1, v_2)$  equal to the (ordered) product of the weights of the arcs on the directed path from  $v_1$  to  $v_2$  if  $v_1 \neq v_2$  and  $D_{(T; \phi; w)}(v_1, v_2)$  equal to  $1_{\mathcal{G}}$  if  $v_1 = v_2$ . Define  $d_{(T; \phi; w)}: X \times X \rightarrow \mathcal{G}$  by setting

$$d_{(T; \phi; w)}(i, j) := D_{(T; \phi; w)}(\phi(i), \phi(j)),$$

for all  $i, j \in X$ . We sometimes drop or abbreviate the subscripts on an  $D_{(T; \phi; w)}$  and  $d_{(T; \phi; w)}$  and write, for example,  $d_T(i, j)$  or even just  $d(i, j)$  if there is no change of ambiguity.

- A *proximity mapping* is any function  $\delta: X \times X \rightarrow \mathcal{G}$  that satisfies  $\delta(i, i) = 1_{\mathcal{G}}$  for all  $i \in X$ . Furthermore, such a mapping is *tree proximity* if there is an  $X$ -tree  $(V, E; \phi)$  with a weight function  $w: \vec{E} \rightarrow \mathcal{G}$  such that, for all  $i, j \in X$ ,  $d_{(T; \phi; w)}(i, j) = \delta(i, j)$ ; in which case  $(T; \phi; w)$  is said to be a *tree representation* of  $\delta$ . If, in addition,  $(T; \phi)$  is a phylogenetic  $X$ -tree and each interior edge is properly weighted, then  $(T; \phi; w)$ , or more briefly  $(T; w)$ , is said to be a *standard tree representation* of  $\delta$ .

Before proving Proposition 2.1, we describe how a tree representation  $(T; \phi; w)$  of a proximity map  $\delta$  gives rise to an associated tree representation  $(T'; \phi'; w')$  of  $\delta$  in which  $(T'; \phi')$  is a phylogenetic  $X$ -tree. For all  $v \in V$  (the set of vertices of  $T$ ), let  $S(v) := \{i \in X : \phi(i) = v\}$  and let  $s(v) := |S(v)|$ . For each interior vertex  $v \in V$  with  $s(v) > 0$  and for each leaf  $v \in V$  with  $s(v) > 1$ , let us make  $v$  the endpoint of  $s(v)$  new edges, and modify  $\phi$  so that, instead of mapping  $S(v)$  to  $v$ , we map  $S(v)$  bijectively to the endpoints of the new edges, thereby creating a phyloge-

netic  $X$ -tree  $(T'; \phi')$ . Let  $w'$  denote the extension of  $w$  to the arcs of  $T'$  by assigning the value  $1_{\mathcal{G}}$  to both arcs of each newly created edge. We will refer to  $(T'; \phi'; w')$  as the *phylogenetic expansion* of  $(T; \phi; w)$ .

**PROPOSITION 2.1.** *Let  $\delta: X \times X \rightarrow \mathcal{G}$  be a tree proximity map. Then there exists a standard tree representation of  $\delta$ .*

*Proof.* By obtaining the phylogenetic expansion of some tree representation of  $\delta$  if necessary, we may assume that we have a tree representation  $(T; \phi; w)$  of  $\delta$  for which  $(T; \phi)$  is a phylogenetic  $X$ -tree with  $T = (V, E)$ . We complete the proof by showing how  $(T; \phi; w)$  can be transformed to a standard tree representation of  $\delta$ .

Suppose that  $u$  and  $v$  are adjacent interior vertices of  $(T; \phi; w)$  with the property that  $w(u, v)w(v, u) = 1_{\mathcal{G}}$ . Let  $(T; \phi; w')$  be obtained from  $(T; \phi; w)$  by replacing  $w$  with the weight function  $w'$  defined, for all distinct  $v_1$  and  $v_2$  of  $V - \{u\}$ , by  $w'(u, v_1) = w(v, u)w(u, v_1)$ ,  $w'(v_1, u) = w(v_1, u)w(v, u)^{-1}$ , and  $w'(v_1, v_2) = w(v_1, v_2)$ , provided the respective pairs are in  $\vec{E}$ . Thus  $w'(u, v) = 1_{\mathcal{G}}$  and  $w'(v, u) = 1_{\mathcal{G}}$ . Using the fact that  $(T; \phi)$  is a phylogenetic  $X$ -tree, a routine check shows that  $(T; \phi; w')$  is a tree representation of  $\delta$ . Let  $T_1$  denote the tree obtained from  $T$  by contracting  $\{u, v\}$  and let  $E_1$  denote the edge set of  $T_1$ . Then the tree  $(T_1; \phi; w'')$ , where the mapping  $w'': \vec{E}_1 \rightarrow \mathcal{G}$  is defined, for all  $(v_1, v_2) \in \vec{E}_1$ , by  $w''(v_1, v_2) = w'(v_1, v_2)$ , is also a tree representation of  $\delta$ . Moreover, it is easily checked that the return-trip weight of every edge in  $(T; \phi; w')$  is equal to  $1_{\mathcal{G}}$  if and only if it is equal to  $1_{\mathcal{G}}$  in  $(T; \phi; w)$ . Hence, in  $(T_1; \phi; w'')$ , the number of properly weighted interior edges is one less than that for  $(T; \phi; w)$ . By continuing this process if necessary, we eventually obtain a standard tree representation of  $\delta$ . ■

*Remark.* In contrast to the classical real-valued symmetric setting, a tree proximity map may not have a tree representation in which the weighting function is proper on *all* edges. In the proof of Theorem 3.1, we outline an explicit polynomial-time construction of a standard tree representation of a tree proximity map.

Before proceeding further, we require the definition of several maps, each of which are essential to the proofs of the main theorems of this paper.

**DEFINITIONS.** • Given a tree  $T = (V, E)$ , a *discriminating  $\mathcal{G}$ -dating map* is a function  $t: V \rightarrow \mathcal{G}$  with the property that if  $\{u, v\}$  is an interior edge of  $T$ , then  $t(u) \neq t(v)$ .

• Given a proximity map  $\delta: X \times X \rightarrow \mathcal{G}$  and an element  $x$  in  $X$ , there is an important associated map  $\delta_x: X \times X \rightarrow \mathcal{G}$  defined, for all

$i, j \in X$ , by

$$\delta_x(i, j) := \delta(x, i) \delta(j, i)^{-1} \delta(j, x).$$

Note that  $\delta_x$  is not usually a proximity map since we generally have  $\delta_x(i, i) \neq 1_{\mathcal{G}}$ . We may regard  $\delta_x$  as the non-Abelian analogue of a classical transformation for real-valued, symmetric proximity maps (described, for example, by Farris, Kluge, and Eckardt [10]).

• Given a phylogenetic  $X$ -tree  $T = (V, E)$ , a discriminating  $\mathcal{G}$ -dating map  $t: V \rightarrow \mathcal{G}$ , an element  $x$  in  $X$ , and a proximity map  $\delta: X \times X \rightarrow \mathcal{G}$ , we describe two associated mappings:

(i) A map  $d_x^{(T;t)}: X \times X \rightarrow \mathcal{G}$  which is defined as follows. For elements  $u$  and  $v$  in  $V$ , write  $u \leq_x v$  if  $u$  lies on the path from  $x$  to  $v$ . For all  $i, j \in X$ , set

$$d_x^{(T;t)}(i, j) := t(\text{glb}_{\leq_x}(i, j)),$$

where  $\text{glb}_{\leq_x}$  denotes the greatest lower bound under the partial order  $\leq_x$ .

(ii) An arc weighting function  $w = w_{t,x}: \vec{E} \rightarrow \mathcal{G}$  which is defined as follows. To each pair of arcs  $(u, v)$  and  $(v, u)$ , assign the weights  $w(u, v)$  and  $w(v, u)$ , respectively, so that:

\* if  $v = i \in X - \{x\}$ , set

$$w(u, i) := t(u)^{-1} t(i) \delta(i, x)^{-1} \quad \text{and} \quad w(i, u) := \delta(i, x);$$

\* otherwise, if  $u \leq_x v$  or  $u = x$ , set

$$w(u, v) := t(u)^{-1} t(v) \quad \text{and} \quad w(v, u) := 1_{\mathcal{G}}.$$

• Lastly, two other mappings are needed. Suppose that  $(T; w)$  is a standard tree representation for a tree proximity  $\delta$  with  $T = (V, E)$ . Let  $x \in X$ .

(i) Define  $t = t^{(T;w;x)}: V \rightarrow \mathcal{G}$  as follows. If  $(u_1, v_1), (u_2, v_2), \dots, (u_k, v_k)$  denotes the arcs on the path from  $x$  to  $v$  (so  $u_1 = x$  and  $v_k = v$ ), then set

$$t(v) := w(u_1, v_1) w(u_2, v_2) \cdots w(u_k, v_k) w(v_k, u_k) \cdots \\ w(v_2, u_2) w(v_1, u_1).$$

In other words, for all  $v \in V$ ,  $t(v)$  is the ordered product of the weights of the arcs on the directed path from  $x$  to  $v$  multiplied by the ordered

product of the weights on the directed path from  $v$  back to  $x$ . Since  $(T; w)$  is a standard tree representation (and so each interior edge is properly weighted), it follows that  $t^{(T; w; x)}$  is a discriminating  $\mathcal{G}$ -dating map.

(ii) The second map  $t_{\delta, x}: V \rightarrow \mathcal{G}$  is defined as follows: for each  $v \in V$ , select elements  $i$  and  $j$  of  $X$  so that  $v$  is the greatest lower bound (under the partial order  $\leq_x$ ) of  $i$  and  $j$ , and set

$$t_{\delta, x}(v) := \delta_x(i, j) = \delta(x, i) \delta(j, i)^{-1} \delta(j, x).$$

That  $t_{\delta, x}$  is well defined (i.e., independent of the choice of  $i$  and  $j$ ) so that, in particular,  $\delta_x(i, j) = \delta_x(j, i)$  (for all  $i, j \in X$ ) whenever  $\delta$  is a tree proximity, and, moreover,  $t_{\delta, x}$  is a discriminating  $\mathcal{G}$ -dating map, follows from the first part of Lemma 2.2.

The following lemma establishes some important relationships between the above mappings.

LEMMA 2.2. *Let  $\delta: X \times X \rightarrow \mathcal{G}$  be a proximity map and let  $x$  be an element of  $X$ .*

1. *If  $(T; w)$  is a standard tree representation of  $\delta$ , then*

$$t_{\delta, x} = t^{(T; w; x)}$$

and

$$\delta_x = d_x^{(T; t')},$$

where  $t' := t^{(T; w; x)}$  ( $= t_{\delta, x}$ ).

2. *Conversely, if  $\delta_x = d_x^{(T; t')}$  for some phylogenetic  $X$ -tree  $T$  and discriminating  $\mathcal{G}$ -dating map  $t$ , then  $(T; w_{t, x})$  is a standard tree representation of  $\delta$ .*

*Proof.* Part 1. To prove the first half of Part 1, let  $v$  be an element of  $V$ , the set of vertices of  $T$ , and choose elements  $i$  and  $j$  of  $X$  so that  $v = \text{glb}_{\leq_x}(i, j)$  in  $T$ . Let  $p_v$  (resp.,  $q_v$ ) be the ordered product of arc weights on the path from  $x$  to  $v$  (resp.,  $v$  to  $x$ ) in  $T$ . Furthermore, let  $p_i$  (resp.,  $q_j$ ) be the ordered product of arc weights on the path from  $v$  to  $i$  (resp.,  $j$  to  $v$ ) in  $T$ . Since  $(T; w)$  is a tree representation of  $\delta$ , it follows that

$$\begin{aligned} t_{\delta, x}(v) &= \delta(x, i) \delta(j, i)^{-1} \delta(j, x) = p_v p_i (q_j p_i)^{-1} q_j q_v \\ &= p_v q_v = t^{(T; w; x)}(v), \end{aligned}$$

as required.

For the second half of Part 1, set  $t' := t_{\delta, x}$ . Since  $t_{\delta, x} = t^{(T; w; x)}$  and since  $t^{(T; w; x)}$  is a discriminating  $\mathcal{G}$ -dating map,  $t_{\delta, x}$  is a discriminating  $\mathcal{G}$ -dating map. Now, for all  $i, j \in X$ , we have

$$d_x^{(T; t')}(i, j) = t_{\delta, x}(\text{glb}_{\leq x}(i, j)) = \delta(x, i)\delta(j, i)^{-1}\delta(j, x) = \delta_x(i, j),$$

as required.

Part 2. Suppose that  $\delta_x = d_x^{(T; t)}$ , for some phylogenetic  $X$ -tree  $T$  and discriminating  $\mathcal{G}$ -dating map  $t$ . First, note that, by the definition of  $w_{t, x}$  and the fact that  $t(u) \neq t(v)$  for each interior edge  $\{u, v\}$  of  $T$ , we see that each interior edge of  $T$  is properly weighted. We complete the proof of Part 2 by verifying that  $d_{(T; w_{t, x})}(i, j) = \delta(i, j)$ , for all  $i, j \in X$ . Let  $I$  denote the cardinality of  $\{x, i, j\}$ . Depending on the value of  $I$ , there are three cases to consider:

- $I = 1$ . In this case,  $d_{(T; w_{t, x})}(x, x) = 1_{\mathcal{G}} = \delta(x, x)$ , as required.

- $I = 2$ . In this case, we may assume that  $i = x$ . Since  $\delta_x = d_x^{(T; t)}$ , we deduce that  $t(x) = 1_{\mathcal{G}}$  and so, by the definition of  $w_{t, x}$ , we have  $d_{(T; w_{t, x})}(x, j) = t(j)\delta(j, x)^{-1}$ . Therefore, as  $t(j) = t(\text{glb}_{\leq x}(j, j)) = d_x^{(T; t)}(j, j) = \delta_x(j, j) = \delta(x, j)\delta(j, x)$ , it follows that  $d_{(T; w_{t, x})}(x, j) = \delta(x, j)$ . Furthermore, from the definition of  $w_{t, x}$ , we directly get  $d_{(T; w_{t, x})}(j, x) = \delta(j, x)$ , as required.

- $I = 3$ . By the definition of  $w_{t, x}$ , we have

$$d_{(T; w_{t, x})}(i, j) = \delta(i, x)t(\text{glb}_{\leq x}(i, j))^{-1}t(j)\delta(j, x)^{-1}.$$

Now

$$\begin{aligned} t(\text{glb}_{\leq x}(i, j)) &= d_x^{(T; t)}(i, j) = d_x^{(T; t)}(j, i) \\ &= \delta_x(j, i) = \delta(x, j)\delta(i, j)^{-1}\delta(i, x), \end{aligned}$$

which also implies that  $t(j) = t(\text{glb}_{\leq x}(j, j)) = \delta(x, j)\delta(j, x)$ . Therefore  $d_{(T; w_{t, x})}(i, j) = \delta(i, j)$ , as required. ■

Combining Proposition 2.1 with the last lemma, we get Corollary 2.3.

**COROLLARY 2.3.** 1. *A proximity map  $\delta: X \times X \rightarrow \mathcal{G}$  is a tree proximity if and only if  $\delta_x = d_x^{(T; t)}$  for some phylogenetic  $X$ -tree  $T$  and discriminating  $\mathcal{G}$ -dating map  $t$ .*

2. *Suppose that  $\delta$  is a tree proximity. If  $T$  is the phylogenetic tree involved in a standard tree representation of  $\delta$ , then  $(T; w_{t, x})$  is a standard tree representation of  $\delta$ , where  $t = t_{\delta, x}$ .*



The next proposition is an immediate consequence of [5, Theorem 2], the main theorem of [5].

**PROPOSITION 2.4.** *Let  $\delta: X \times X \rightarrow \mathcal{G}$  be a proximity map and let  $x$  be an element of  $X$ . Then there exists a phylogenetic  $X$ -tree  $T$  and a discriminating  $\mathcal{G}$ -dating map  $t$  such that  $\delta_x = d_x^{(T;t)}$  if and only if  $\delta_x$  satisfies the following conditions:*

- (U1)  $\delta_x(i, j) = \delta_x(j, i)$ , for all  $i, j \in X$ ;
- (U2)  $|\{\delta_x(i, j), \delta_x(i, k), \delta_x(j, k)\}| \leq 2$ , for all  $i, j, k \in X$ ; and
- (U3) *there exist no pairwise distinct elements  $i, j, k$ , and  $l$  of  $X$  with*  

$$\delta_x(i, j) = \delta_x(j, k) = \delta_x(k, l) \neq \delta_x(j, l) = \delta_x(l, i) = \delta_x(i, k).$$

Furthermore, up to canonical isomorphism,  $T$  is unique.

Combining Corollary 2.3 with Proposition 2.4, we get Corollary 2.5.

**COROLLARY 2.5.** *A proximity map  $\delta: X \times X \rightarrow \mathcal{G}$  is a tree proximity map if and only if  $\delta_x$  satisfies (U1), (U2), and (U3) for some  $x \in X$ .*

The existence part of Theorem 3.1 is proved via Corollary 2.5.

### 3. MAIN RESULTS

We are now ready to state and prove our two main (existence and uniqueness) results, Theorems 3.1 and 3.2. Note that an explanation for the slight complication concerning “ $H_\delta$ ” in Part 2 of the statement of Theorem 3.1 is given in the remark immediately following the proof of Theorem 3.1.

**THEOREM 3.1.** *Let  $\delta: X \times X \rightarrow \mathcal{G}$  be a proximity map. Let  $H_\delta$  denote the following (finite) subset of  $\mathcal{G}$ :*

$$\{\delta(i, k)\delta(j, k)^{-1}\delta(j, l)\delta(i, l)^{-1} : i, j, k, l \in X\}.$$

1. *If  $\delta$  is a tree proximity map, then  $\delta$  satisfies the following conditions:*

(P1) *For all distinct elements  $i, j$ , and  $k$  of  $X$ ,*

$$\delta(i, j)\delta(k, j)^{-1}\delta(k, i) = \delta(i, k)\delta(j, k)^{-1}\delta(j, i).$$

(P2) *For all four distinct elements of  $X$ , we can order these elements as  $i, j, k$ , and  $l$  so that*

$$\delta(i, k)\delta(j, k)^{-1} = \delta(i, l)\delta(j, l)^{-1}.$$

2. If  $\delta$  satisfies both (P1) and (P2), and  $H_\delta$  has no elements of order 2, then  $\delta$  is a tree proximity map.

Furthermore, a standard tree representation of  $\delta$ , if one exists, can be constructed in polynomial time from  $\delta$ .

*Proof.* If  $\delta$  is a tree proximity map, then it is straightforward to check that (P1) and (P2) must hold by cancelling the products of the appropriate arc weights in  $\mathcal{G}$ . Thus part 1 of Theorem 3.1 holds.

Before proving Part 2 of Theorem 3.1, we note the following. If  $\delta(i, k)\delta(j, k)^{-1} = \delta(i, l)\delta(j, l)^{-1}$ , for some elements  $i, j, k$ , and  $l$  of  $X$ , then it is easily checked using (P1) that

$$\delta(k, i)\delta(l, i)^{-1} = \delta(k, j)\delta(l, j)^{-1}$$

also holds. We freely use this observation in the proof that follows.

Let  $x$  be an element of  $X$ . To prove Part 2, it suffices to show, by Corollary 2.5, that  $\delta_x$  satisfies conditions (U1), (U2), and (U3) as listed in the statement of Proposition 2.4. We now show that this is indeed the case.

For all  $i, j \in X$ , (P1) shows that  $\delta_x$  satisfies (U1). Furthermore, for all  $i, j, k \in X$ , (P2) together with (U1) shows that  $\delta_x$  satisfies (U2). The proof that (U3) holds for  $\delta_x$  is

3.1.1.  $\delta_x$  satisfies (U3).

*Proof.* Suppose that  $i, j, k$ , and  $l$  are pairwise distinct elements of  $X$  with

$$\delta_x(i, j) = \delta_x(j, k) = \delta_x(k, l),$$

and

$$\delta_x(j, l) = \delta_x(l, i) = \delta_x(i, k).$$

We prove (3.1.1) by showing that  $\delta_x(i, j)$ ,  $\delta_x(j, k)$ ,  $\delta_x(k, l)$ ,  $\delta_x(j, l)$ ,  $\delta_x(l, i)$ , and  $\delta_x(i, k)$  are all equal.

If  $|\{x, i, j, k, l\}| = 4$ , then it is clear that (3.1.1) holds. Therefore assume that  $|\{x, i, j, k, l\}| = 5$ . Depending on the relationship between  $i, j, k$ , and  $l$  given by (P2) and noting the observation above, there are three cases to consider:

- (i)  $\delta(i, k)\delta(j, k)^{-1} = \delta(i, l)\delta(j, l)^{-1}$ ;
- (ii)  $\delta(i, j)\delta(k, j)^{-1} = \delta(i, l)\delta(k, l)^{-1}$ ; and
- (iii)  $\delta(i, j)\delta(l, j)^{-1} = \delta(i, k)\delta(l, k)^{-1}$ .

We denote the above equations as (i)–(iii), respectively. Moreover, in the analysis of Cases (i)–(iii), we freely use the fact that (U1) holds for  $\delta_x$ .

*Case (i).* Since  $\delta_x(l, i) = \delta_x(k, i)$ , we have  $\delta(x, l)\delta(i, l)^{-1} = \delta(x, k)\delta(i, k)^{-1}$ , which implies that

$$\delta(x, l)\delta(i, l)^{-1}\delta(i, k) = \delta(x, k). \quad (1)$$

Now, by (i),  $\delta(j, l)^{-1} = \delta(i, l)^{-1}\delta(i, k)\delta(j, k)^{-1}$  and so

$$\begin{aligned} \delta_x(j, l) &= \delta_x(l, j) = \delta(x, l)\delta(j, l)^{-1}\delta(j, x) \\ &= \delta(x, l)\delta(i, l)^{-1}\delta(i, k)\delta(j, k)^{-1}\delta(j, x) \\ &= \delta(x, k)\delta(j, k)^{-1}\delta(j, x), \quad \text{by (1),} \\ &= \delta_x(k, j) = \delta_x(j, k), \end{aligned}$$

completing the proof of (U3) for Case (i).

*Case (ii).* The proof of (U3) for Case (ii) is analogous to that of Case (i). We omit the details and just note that we first deduce  $\delta(x, k)\delta(j, k)^{-1}\delta(j, i) = \delta(x, i)$  via the fact that  $\delta_x(i, j) = \delta_x(k, j)$ , and then show  $\delta_x(k, l) = \delta_x(l, i)$ .

*Case (iii).* Since  $\delta_x(j, i) = \delta_x(k, l)$  and since  $\delta_x(j, l) = \delta_x(k, i)$ , we have

$$\delta(x, j)\delta(i, j)^{-1}\delta(i, x) = \delta(x, k)\delta(l, k)^{-1}\delta(l, x), \quad (2)$$

and

$$\delta(x, j)\delta(l, j)^{-1}\delta(l, x) = \delta(x, k)\delta(i, k)^{-1}\delta(i, x), \quad (3)$$

respectively. By combining (2) and (3), we deduce that

$$\delta(l, k)^{-1}\delta(l, x)\delta(i, x)^{-1}\delta(i, j) = \delta(i, k)^{-1}\delta(i, x)\delta(l, x)^{-1}\delta(l, j),$$

which in turn implies that

$$\delta(i, k)\delta(l, k)^{-1}\delta(l, x)\delta(i, x)^{-1} = \delta(i, x)\delta(l, x)^{-1}\delta(l, j)\delta(i, j)^{-1}. \quad (4)$$

Substituting (iii) into (4), we get

$$\delta(i, j)\delta(l, j)^{-1}\delta(l, x)\delta(i, x)^{-1} = \delta(i, x)\delta(l, x)^{-1}\delta(l, j)\delta(i, j)^{-1}.$$

Since  $H_\delta$  has no elements of order 2, the last equation implies that

$$\delta(i, x)\delta(l, x)^{-1}\delta(l, j)\delta(i, j)^{-1} = 1_{\mathcal{G}}. \quad (5)$$

Having established (5), we complete the proof of (U3) for Case (iii) as follows. By (iii),  $\delta(i, k)^{-1} = \delta(l, k)^{-1}\delta(l, j)\delta(i, j)^{-1}$ , and so

$$\begin{aligned}\delta_x(i, k) &= \delta_x(k, i) = \delta(x, k)\delta(i, k)^{-1}\delta(i, x) \\ &= \delta(x, k)\delta(l, k)^{-1}\delta(l, j)\delta(i, j)^{-1}\delta(i, x) \\ &= \delta(x, k)\delta(l, k)^{-1}\delta(l, x), \quad \text{by (5),} \\ &= \delta_x(k, l),\end{aligned}$$

as required.

We conclude that  $\delta_x$  satisfies (U1)–(U3) and so, by Corollary 2.5,  $\delta$  is a tree proximity map and this completes the proof of Part 2.

Lastly, we describe a polynomial time algorithm for finding a standard tree representation of  $\delta$ . First, we provide a construction of a phylogenetic  $X$ -tree that turns out to be isomorphic to the underlying phylogenetic  $X$ -tree of a standard tree representation of  $\delta$ . For a tree proximity map  $\delta$  and an element  $x$  in  $X$ , let  $R(\delta, x)$  denote the set of  $x$ -rooted phylogenetic trees (that is, trees rooted on leaf  $x$ ) which is constructed as follows. For each pairwise disjoint triple  $i, j, k \in X$ , if

$$\delta_x(i, j) \neq \delta_x(i, k) = \delta_x(j, k),$$

then place the  $x$ -rooted tree  $ij|kx$ , as shown in Fig. 1, into  $R(\delta, x)$ . Let  $A[R(\delta, x)]$  denote the  $x$ -rooted tree constructed from  $R(\delta, x)$  by applying the algorithm of Aho *et al.* [1] (see also [6, 13]). Briefly, in this algorithm, one first constructs a graph  $G$  having vertex set  $X - \{x\}$  and with an edge between any two vertices  $i$  and  $j$  precisely if there exists  $k \in X - \{x\}$  such that  $ij|kx \in R(\delta, x)$ . One then takes the connected components of this graph, which form the top “clusters” of the tree, and continues this process recursively on the vertices of each component. For further details see [6, 13].

We now show that if  $(T; w)$  is a standard tree representation of  $\delta$ , then

$$A[R(\delta, x)] \equiv T. \quad (6)$$

To prove (6), we argue by induction based on the number of interior vertices in the longest path of  $T$  that starts at  $x$ , when one considers  $T$  as an  $x$ -rooted tree. Let  $h(T)$  denote this number. If  $h(T) = 1$ , then, as  $T$  is part of a standard tree representation of  $\delta$ , it follows by the first part of Lemma 2.2 that  $R(\delta, x)$  is empty and so (6) holds.

Now assume that  $h(T) > 1$  and that (6) holds for all trees in a standard tree representation of  $\delta$  with fewer interior vertices in the longest path starting at  $x$ . For  $r > 1$ , let  $V_1, V_2, \dots, V_r$  denote the vertex sets of the

subtrees of  $T$ , other than the isolated vertex  $x$ , incident with the vertex of  $T$  adjacent to  $x$ . For all  $p \in \{1, 2, \dots, r\}$ , let  $X_p := \phi^{-1}(V_p)$ . Thus  $X_1, X_2, \dots, X_r$  forms a partition of  $X - \{x\}$ .

Let  $G$  be the graph described above in the brief description of the algorithm. To prove the induction step of the proof, it suffices to show that  $X_1, X_2, \dots, X_r$  are precisely the vertex sets of the connected components of  $G$ . That is, for some  $p \in \{1, 2, \dots, r\}$ , elements  $i$  and  $j$  are both in  $X_p$  if and only if there exists  $k \in X - \{x\}$  such that  $ij|kx \in R(\delta, x)$ . We now show that this is indeed the case.

Suppose that  $i, j \in X_p$ , for some  $p \in \{1, 2, \dots, r\}$ . Let  $v = \text{glb}_{\leq_x}(i, j)$  in  $T$ . Since  $i, j \in X_p$ , there exists an interior vertex  $u$  in  $T$  and a  $k$  in  $X$  such that, in  $T$ , vertices  $u$  and  $v$  are adjacent and  $u = \text{glb}_{\leq_x}(i, k) = \text{glb}_{\leq_x}(j, k)$ . By Part 1 of Lemma 2.2, the map  $t_{\delta, x}: V \rightarrow \mathcal{G}$ , where  $V$  denotes the vertex set of  $T$ , is a discriminating  $\mathcal{G}$ -dating map and so  $t_{\delta, x}(u) \neq t_{\delta, x}(v)$ . Therefore

$$d_x^{(T:t)}(i, j) \neq d_x^{(T:t)}(i, k) = d_x^{(T:t)}(j, k),$$

where  $t = t_{\delta, x}$ . By Part 1 of Lemma 2.2,  $\delta_x = d_x^{(T:t)}$  and so  $ij|kx \in R(\delta, x)$ .

Now suppose that there is no  $p$  in  $\{1, 2, \dots, r\}$  such that  $i, j \in X_p$ . An argument similar to that used in the last paragraph, shows that there is no  $k \in X - \{x\}$  such that  $ij|kx \in R(\delta, x)$ . This establishes (6).

Having established (6), we see that  $A[R(\delta, x)]$  is part of a standard tree representation of  $\delta$ . By Part 2 of Corollary 2.3, the pair  $(A[R(\delta, x)]; w_{t, x})$ , for  $t = t_{\delta, x}$ , provides a standard tree representation of  $\delta$ . Furthermore, the tree  $A[R(\delta, x)]$  can be constructed in polynomial time (see [1, 6, 13]), and once this tree is constructed, the arc function  $w_{t, x}$  can also be constructed in polynomial time. This completes the proof of Theorem 3.1.

■

*Remark.* The condition on  $H_\delta$  in the statement of Theorem 3.1 is necessary as there exists a group  $\mathcal{G}$  with elements of order 2 and a proximity map  $\delta: X \times X \rightarrow \mathcal{G}$  such that (P1) and (P2) are satisfied, but in which there is no tree representation of  $\delta$ . An example is provided by the construction in [4] used to illustrate the “necessary part” of [4, Proposition 1(2)].

Given a standard tree representation  $(T; w)$  of a tree proximity map  $\delta$  our second main result shows that, up to isomorphism,  $T$  is determined by  $\delta$ , and the arc weighting  $w: \vec{E} \rightarrow \mathcal{G}$  is partially determined. More precisely, although  $w$  is not completely determined (as pointed out by [2, 8, 11]), the return-trip weights of every exterior edge as well as, up to conjugacy, the return-trip weights of every interior edge of  $(T; w)$  can be obtained (this was established for the particular group analyzed in [2]).

Moreover, we show that the arc weights can be arbitrarily specified on a certain subset of arcs, but once this is done, then all the remaining arc weights are determined.

Before stating Theorem 3.2, we note the following. If  $T$  and  $T'$  are two isomorphic trees, then one can identify the set of vertices (resp., edges) of  $T'$  as being equal to the set of vertices (resp., edges) of  $T$ . For the sake of simplicity and without ambiguity, we shall treat the vertices (resp., edges) of two such trees in the statement and proof of Theorem 3.2 as equivalent.

**THEOREM 3.2.** *Let  $\delta: X \times X \rightarrow \mathcal{G}$  be a tree proximity map. Suppose that  $(T; w)$  and  $(T'; w')$  are both standard tree representations of  $\delta$ . Then:*

1.  $T$  is isomorphic to  $T'$ .
2. Let  $e = \{u, v\}$  be an edge of  $T$ .
  - (i) If  $e$  is an exterior edge, then

$$w(u, v)w(v, u) = w'(u, v)w'(v, u).$$

- (ii) If  $e$  is an interior edge, then

$$w(u, v)w(v, u) \cong w'(u, v)w'(v, u),$$

where  $\alpha \cong \beta$  denotes conjugacy in  $\mathcal{G}$ , that is, there exists an element  $\gamma$  in  $\mathcal{G}$  such that  $\alpha = \gamma\beta\gamma^{-1}$ .

3. Select an interior (resp., exterior) edge,  $\{u, v\}$ , say, of  $T$ . For all  $\xi, \zeta \in \mathcal{G}$  such that  $\xi\zeta \cong w(u, v)w(v, u)$  (resp.,  $\xi\zeta = w(u, v)w(v, u)$ ), there exists a standard tree representation  $(T; w'')$  with

$$w''(u, v) = \xi \quad \text{and} \quad w''(v, u) = \zeta.$$

4. Let  $\overset{\circ}{E} \subset E$  denote the set of interior edges of  $T$ . Then there exists a subset  $A$  of  $\overset{\circ}{E}$ , with  $|A| = 1 + |\overset{\circ}{E}|$ , such that

$$w|_A = w'|_A \quad \Rightarrow \quad w = w'.$$

Furthermore, provided  $|\overset{\circ}{E}| \geq 1$ , one can extend an arbitrary assignment of elements of  $\mathcal{G}$  to the members of  $A$  to a weight function from  $\overset{\circ}{E}$  into  $\mathcal{G}$  which, together with  $T$ , gives a standard tree representation of  $\delta$ . Moreover, all standard tree representations of  $\delta$  can be obtained in this way.

*Proof.* Part 1. Equation (6) shows that  $T$  is determined by  $\delta$ , and provides, moreover, a polynomial time constructive algorithm. Alternatively, the result may be deduced from Proposition 2.4 as follows. From the proof of Theorem 3.1,  $\delta_x$  satisfies (U1)–(U3). Therefore, by combining the first part of Lemma 2.2 with Proposition 2.4, we deduce that  $T$  is isomorphic to  $T'$ .

Part 2. Here we freely use the fact, from the previous paper, that  $T$  is isomorphic to  $T'$ .

To prove (i), suppose that  $e = \{i, u\}$  is an exterior edge of  $T$ , where  $i \in X$ . If  $i$  and  $u$  are the only vertices of  $T$ , then (i) holds. Therefore assume that  $T$  has at least three vertices. Let  $j$  and  $k$  be elements of  $X - \{i\}$  such that the path from  $j$  to  $k$  in  $T$  is incident with  $u$ . It follows that

$$\delta(i, j) \delta(k, j)^{-1} \delta(k, i) = w(i, u) w(u, i),$$

completing the proof of (i).

To prove (ii), suppose that  $e = \{u, v\}$  is an exterior edge of  $T$ . Now let  $i$  and  $j$  be elements of  $X$  such that the path from  $i$  to  $j$  is incident with  $u$ , but not with  $v$ . Similarly, let  $k$  and  $l$  be elements of  $X$  such that the path from  $k$  to  $l$  is incident with  $v$ , but not with  $u$ . Then

$$\begin{aligned} & \delta(i, l) \delta(k, l)^{-1} \delta(k, j) \delta(i, j)^{-1} \\ &= D_{(T; w)}(i, u) w(u, v) w(v, u) D_{(T; w)}(i, u)^{-1}, \end{aligned}$$

and

$$\begin{aligned} & \delta(i, l) \delta(k, l)^{-1} \delta(k, j) \delta(i, j)^{-1} \\ &= D_{(T'; w')}(i, u) w'(u, v) w'(v, u) D_{(T'; w')}(i, u)^{-1}. \end{aligned}$$

By equating the right-hand sides of the last two equations, and then multiplying the resulting equation on the left by  $D_{(T; w)}(i, u)^{-1}$  and on the right by  $D_{(T; w)}(i, u)^{-1}$ , we get the desired result. This completes the proof of (ii).

Part 3. Suppose that  $\{u, v\}$  is an interior edge of  $T$ . Let  $\alpha = w(u, v)$  and  $\beta = w(v, u)$ , and suppose that  $\xi\zeta \cong \alpha\beta$ , that is,  $\xi\zeta = \gamma\alpha\beta\gamma^{-1}$  for some  $\gamma \in \mathcal{G}$ . Let  $\{u_1, u_2, \dots, u_r\}$  be the set of vertices in  $T$  adjacent  $u$  other than  $v$ , and let  $\{v_1, v_2, \dots, v_s\}$  be the set of vertices in  $T$  adjacent to  $v$  other than  $u$ . Let  $w'' : \vec{E} \rightarrow \mathcal{G}$  denote the arc weighting function defined as

- $w''(u, v) := \xi$  and  $w''(v, u) := \zeta$ ;
- $w''(u, u_i) := \gamma w(u, u_i)$  and  $w''(u_i, u) := w(u_i, u) \gamma^{-1}$ , for all  $i \in \{1, 2, \dots, r\}$ ;
- $w''(v, v_j) := \xi^{-1} \gamma \alpha w(v, v_j)$  and  $w''(v_j, v) := w(v_j, v) \beta \gamma^{-1} \zeta^{-1}$ , for all  $j \in \{1, 2, \dots, s\}$ ; and
- $w''$  agrees with  $w$  on all other arcs.

It is easily checked that each edge of  $T$  is properly weighted under the arc weighting  $w''$ . Furthermore, for all  $i, j \in X$ , a case analysis (depending on which of the above arcs are crossed in the path from  $i$  to  $j$ ) using elementary cancellation of products in the group  $\mathcal{G}$  shows that  $\delta_{(T; w'')}(i, j) = \delta_{(T; w)}(i, j)$ , as required.

Now suppose that  $\{x, u\}$  is a exterior edge of  $T$ , with  $x \in X$ ,  $w(x, u) = \alpha$ , and  $w(u, x) = \beta$ . Suppose that  $\xi\zeta = \alpha\beta$  and let  $\{u_1, u_2, \dots, u_r\}$  be the set of vertices in  $T$  adjacent to  $u$  other than  $x$ . Let  $w'' : \vec{E} \rightarrow \mathcal{G}$  denote the arc weighting function defined as

- $w''(x, u) := \xi$  and  $w''(u, x) := \zeta$ ;
- $w''(u, u_i) := \xi^{-1}\alpha w(u, u_i)$  and  $w''(u_i, u) := w(u_i, u)\beta\zeta^{-1}$ , for every  $i \in \{1, \dots, r\}$ ; and
- $w''$  agrees with  $w$  on all other arcs.

Again it is easily checked that each edge of  $T$  is properly weighted under the new arc weighting  $w''$ , and that, for all  $i, j \in X$ ,  $\delta_{(T; w'')}(i, j) = \delta_{(T; w)}(i, j)$ , as required.

Part 4. We begin the proof of Part 4 by constructing the desired subset of  $\vec{E}$ . Select an element,  $x$  say, of  $X$ . Let  $e$  be the edge of  $T$  incident with  $x$ . Set  $A$  to be a subset of  $\vec{E}$  such that  $(u, v) \in A$  if and only if  $(v, u) \notin A$  and  $\{u, v\} \in \mathring{E} \cup \{e\}$ . We now show that  $A$  has the properties claimed in the statement of Part 4.

First, to each member of  $A$  assign an arbitrary element of  $\mathcal{G}$ . Let  $w^* : \vec{E} \rightarrow \mathcal{G}$  denote the arc weight function that extends this arbitrary assignment of elements of  $\mathcal{G}$  to the members of  $\vec{E}$  and is constructed as

- For each arc  $(u, v)$  in  $A$ , set  $w^*(v, u)$  so that, if  $u \leq_x v$  (resp.,  $v \leq_x u$ ) holds, the ordered product of the weights of the arcs from  $x$  to  $v$  (resp.,  $u$ ) and back to  $x$  is equal to  $t_{\delta, x}(v)$  (resp.,  $t_{\delta, x}(u)$ ). It is not difficult to see that this can be done recursively (and furthermore uniquely) based on the number of edges separating  $v$  (resp.,  $u$ ) from  $x$ .

- For the remaining arcs in  $\vec{E}$ , if  $v = i \in X$ , then set  $w^*(u, v) := p^{-1}\delta(x, i)$  and  $w^*(v, u) := \delta(i, x)q^{-1}$ , where  $p$  is the ordered product of the arc weights from  $x$  to  $u$  under  $w^*$  and  $q$  is the ordered product of the arc weights from  $u$  to  $x$  under  $w^*$ .

Note that  $w^*$  is well defined. We next show that  $(T; w^*)$  is a tree representation of  $\delta$  by showing that, for all  $i, j \in X$ ,  $d_{(T; w^*)}(i, j) = d_{(T; w)}(i, j)$ .

Set  $d = d_{(T; w)}$ . Clearly, we have  $d_{(T; w^*)}(x, i) = \delta(x, i) = d(x, i)$  and, similarly,  $d_{(T; w^*)}(i, x) = d(i, x)$ , for all  $i \in X$ . So assume that  $i, j$ , and  $x$  are pairwise distinct. Let  $v = \text{glb}_{\leq_x}(i, j)$ , and let  $p_v$  and  $q_v$  be the ordered



products of arc weights from  $x$  to  $v$  and from  $v$  to  $x$ , respectively, in  $(T; w^*)$ . Furthermore, let  $q_i$  and  $p_j$  be the ordered products of arc weights from  $i$  to  $v$  and from  $v$  to  $j$ , respectively, in  $(T; w^*)$ . By the definition of  $w^*$ , we have

$$p_v q_v = t_{\delta, x}(v) = \delta(x, j) \delta(i, j)^{-1} \delta(i, x) = d(x, j) d(i, j)^{-1} d(i, x).$$

Therefore

$$\begin{aligned} d(i, j) &= \left[ d(x, j)^{-1} p_v q_v d(i, x)^{-1} \right]^{-1} \\ &= \left[ (p_v p_j)^{-1} p_v q_v (q_i q_v)^{-1} \right]^{-1}, \\ &\quad \text{since } d = d_{(T; w^*)} \text{ when } x \in \{i, j\}, \\ &= q_i p_j = d_{(T; w^*)}(i, j). \end{aligned}$$

Hence  $(T; w^*)$  is a tree representation of  $\delta$ . Furthermore,  $(T; w^*)$  must be a standard tree representation of  $\delta$ , for otherwise, the phylogenetic  $X$ -tree associated with the standard tree representation of  $\delta$  obtained from  $(T; w^*)$ , by the method described in Proposition 2.1, has fewer internal vertices than  $T$ , contradicting Part 1.

Since  $w^*_{|_A}$  and  $\delta$  determines the weight of each arc under  $w^*$  and since  $T$  is isomorphic to  $T'$ , it follows that all standard tree representations of  $\delta$  can be obtained in this way by making the appropriate assignment of elements of  $\mathcal{S}$  to the members of  $A$  and that if  $w^*_{|_A} = w'_{|_A}$ , then  $w = w'$ . This completes the proof of Part 4 and so Theorem 3.2 is proved. ■

We complete this section of the paper by showing that the main theorem of [4] can be deduced from Theorems 3.1 and 3.2.

Suppose  $\mathcal{S}$  is an Abelian semigroup with identity (we denote the binary operation by addition  $+$ , the identity by  $0$ , and write  $2x$  as shorthand for  $x + x$ ). In [4], the authors considered two further conditions on  $\mathcal{S}$ , namely, cancellation ( $x + y = x + z \Rightarrow y = z$ ) and uniqueness of halves ( $2x = 2y \Rightarrow x = y$ ). These two conditions are easily seen to be equivalent to the condition that  $\mathcal{S}$  embeds in an Abelian group  $\mathcal{G}$  that has no elements of order 2. Thus the following corollary immediately gives the main theorem of [4].

**COROLLARY 3.3.** *Suppose  $\mathcal{G}$  is an Abelian group, with no elements of order 2, and  $\mathcal{S} \subseteq \mathcal{G}$  forms a semigroup. Suppose further that  $\delta: X \times X \rightarrow \mathcal{S}$  is symmetric (i.e.,  $\delta(i, j) = \delta(j, i)$ , for all  $i, j \in X$ ). Then  $\delta$  can be realized*

by a symmetric edge weighting  $w: \vec{E} \rightarrow \mathcal{S}$  of an  $X$ -tree  $(T; \phi)$  if and only if the following four point condition applies:

For all (not necessarily distinct) four points in  $X$ , there exists an ordering of these points,  $i, j, k$ , and  $l$  say, and an element  $\xi$  in  $\mathcal{S}$  such that

$$\delta(i, j) + \delta(k, l) + 2\xi = \delta(i, k) + \delta(j, l) = \delta(i, l) + \delta(j, k). \quad (7)$$

Furthermore, the triple  $(T; \phi; w)$  is uniquely determined by  $\delta$ , provided we insist that no arc of  $T$  has weighting zero.

*Proof.* Regarding the existence of a tree representation of  $\delta$  the ‘‘only if’’ direction is clear. For the ‘‘if’’ part, we note that conditions (P1) and (P2) in the statement of Theorem 3.1 clearly apply, and thus, by Theorem 3.1, there exists a tree representation  $(T; \phi; w')$  of  $\delta$ , where  $(T; \phi)$  is a phylogenetic  $X$ -tree and  $w': \vec{E} \rightarrow \mathcal{G}$ . We wish to show that  $w'$  can be replaced by a function  $w$  that (i) maps into  $\mathcal{S}$  and (ii) is symmetric. To this end, we first establish the following claim: For each edge  $e = \{u, v\}$  of  $T$ , there exists  $\xi \in \mathcal{S}$  such that

$$w'(u, v) + w'(v, u) = 2\xi. \quad (8)$$

To establish (8), there are two cases to consider depending upon  $e$  being either an interior edge of  $T$  or an exterior edge of  $T$ . We consider just the former, since the proof of the latter is similar. Select  $i', j', k', l' \in X$  in such a way that in  $T$  the path between leaves  $i'$  and  $j'$  is incident with  $u$ , but not with  $v$ , while the path between  $k'$  and  $l'$  is incident with  $v$ , but not with  $u$ . Then, as  $G$  is Abelian and denoting  $d_{(T; \phi; w')}$  as  $d'$ , we get

$$d'(i', j') + d'(l', k') + w'(u, v) + w'(v, u) = d'(i', k') + d'(l', j').$$

Since  $d' \equiv \delta$ , the condition described by Eq. (7) (plus the symmetry of  $\delta$ ) guarantees the existence of  $\xi \in \mathcal{S}$  such that

$$d'(i', j') + d'(l', k') + 2\xi = d'(i', k') + d'(l', j'),$$

which in view of the previous equation implies that  $w'(u, v) + w'(v, u) = 2\xi$ , as required to establish the claim.

Now, referring to Eq. (8), set  $w(u, v)$  and  $w(v, u)$  both equal to  $\xi$ , for each edge  $\{u, v\}$  of  $T$ . Let  $d := d_{(T; \phi; w)}$ . Then, for each  $i, j \in X$ ,

$$\begin{aligned} 2d(i, j) &= d(i, j) + d(j, i) = d'(i, j) + d'(j, i) \\ &= \delta(i, j) + \delta(j, i) = 2\delta(i, j), \end{aligned}$$

and so  $2[d(i, j) - \delta(i, j)] = 0$ . Since  $\mathcal{G}$  has no elements of order 2, it follows that  $d \equiv \delta$ , and so  $(T; \phi; w)$  provides the desired tree representation of  $\delta$ .

Regarding the uniqueness of the tree representation, suppose that  $(T_1; \phi_1; w_1)$  and  $(T_2; \phi_2; w_2)$  both provide tree representations of  $\delta$ , where  $w_1$  and  $w_2$  are both symmetric functions taking values in  $\mathcal{S} - \{0\}$ .

For all  $i \in \{1, 2\}$ , let  $(T'_i; w'_i)$  denote the phylogenetic expansion of the tree representation  $(T_i; \phi_i; w_i)$ . Then  $(T'_1; w'_1)$  and  $(T'_2; w'_2)$  are both standard tree representations of  $\delta$ . Consequently, by Theorem 3.2,  $T'_1$  and  $T'_2$  are isomorphic. Therefore, by noting that, for an Abelian group, two elements are conjugates precisely if they are identical, the second part of Theorem 3.2 shows that  $w'_1(u, v) + w'_1(v, u) = w'_2(u, v) + w'_2(v, u)$  for each (isometrically equivalent) edge  $\{u, v\}$  in  $T_1$  and  $T_2$ . But, since  $w'_1$  and  $w'_2$  are both symmetric, this implies that  $2[w'_1(u, v) - w'_2(u, v)] = 0$  and, since  $\mathcal{S}$  has no elements of order 2, this in turn implies that  $w'_1(u, v) = w'_2(u, v)$ . In particular,  $w'_1(u, v) = 0$  precisely if  $w'_2(u, v) = 0$ , which together with the isomorphism between  $T'_1$  and  $T'_2$  and the way in which these trees were constructed from  $(T_1; \phi_1)$  and  $(T_2; \phi_2)$  implies that  $(T_1; \phi_1)$  and  $(T_2; \phi_2)$  are isomorphic, and  $w_1 \equiv w_2$  as required. ■

#### 4. REMARKS

- We return to the problem that motivated our analysis, namely, the case where  $\mathcal{S}$  is the group of  $r \times r$  non-singular real matrices and we have a general  $r$ -state Markov process on a phylogenetic  $X$ -tree  $T$  ([2, 8, 12, 15]). In this setting, each arc  $(u, v)$  of  $T$  has an associated transition matrix  $P^{(u, v)}$  (i.e., a matrix with non-negative entries, and with each row summing to (1)). Assigning arc weights by setting  $w(u, v) := P^{(u, v)}$ , the induced tree proximity value  $\delta(i, j)$  is the net transition matrix of the states at  $j$  conditional on the states at  $i$ . For technical reasons, the following mild restrictions are usually imposed in this model: (i) each transition matrix is neither a singular matrix nor a permutation matrix (this is equivalent to requiring that the matrix determinant is not in  $\{0, \pm 1\}$ ); and (ii), for some vertex in the tree, each state occurs with strictly positive probability (this ensures that some quantities described below are well defined). Restriction (i) clearly implies that  $P^{(u, v)}P^{(v, u)}$  is not the identity matrix for any edge  $\{u, v\}$  of  $T$ . Consequently  $(T; w)$  provides a standard tree representation of  $\delta$ . Now, by elementary probability theory,  $\delta(i, j) = (J^{ii})^{-1}J^{ij}$  where, for  $k, l \in \{i, j\}$ ,  $J^{kl}$  is the  $r \times r$  matrix consisting of the joint probabilities of the states at leaves  $k$  and  $l$ . Thus, by Theorem 3.2, we see immediately that the joint probability distributions of states at pairs of leaves determine  $T$  up to isomorphism (under restrictions (i) and (ii)). This result was established using a less direct approach in [8, 12, 15], essentially by reducing the problem to the classical setting (symmetric edge weightings over  $\mathbb{R}^{\geq 0}$ ).

For completeness, we also sketch this line of argument here. Consider the group homomorphism  $f$  from this particular group  $\mathcal{G}$  into the Abelian group  $\mathbb{R}$  of real numbers under addition, defined by

$$f(M) = -\log(|\det(M)|),$$

where “det” refers to matrix determinant. For any transition matrix  $M$ ,  $\det(M) \in [-1, 1]$  and so  $f$  maps the semigroup of non-singular transition matrices into the semigroup  $\mathbb{R}^{\geq 0}$  of non-negative real numbers under addition. Thus, if we set

$$\delta'(i, j) := \frac{1}{2}(f(\delta(i, j)) + f(\delta(j, i))),$$

then  $\delta'$  is a symmetric tree proximity with a representation on  $T$  via the symmetric arc weight function  $w'$  defined by

$$w'(u, v) = w'(v, u) := \frac{1}{2}(f(w(u, v)) + f(w(v, u))).$$

From assumption (i),  $f$  is strictly positive on  $w(u, v)$  and  $w(v, u)$ , and so  $w'(u, v) = w'(v, u) > 0$  for all edges  $\{u, v\}$  of  $T$ . Thus, by the previous corollary (or indeed the classical result),  $\delta'$  uniquely determines  $T$ . Furthermore, by routine manipulation,

$$\delta'(i, j) = -\log\left(\frac{|\det(J^{ij})|}{\sqrt{\det(J^{ii})\det(J^{jj})}}\right).$$

Thus, under restrictions (i) and (ii), the joint probability distributions of states at pairs of leaves of  $T$  determine  $T$  up to isomorphism via the proximity map  $\delta'$  ([8, 12]). In fact, the slightly simpler proximity map  $\delta''(i, j) := -\log(|\det(J^{ij})|)$ , for  $i \neq j$ , also determines  $T$  up to isomorphism [15].

- Our other main result, Theorem 3.1, also pertains to this particular setting. Conditions (P1) and (P2) translate into a collection of polynomial function identities between the entries of the  $J^{kl}$  matrices—such functions are examples of “phylogenetic invariants” [9] and we note that the invariants described by (P1) are independent of the underlying tree  $T$  (so-called “model invariants”). To date, most investigation of phylogenetic invariants has been for submodels of this general model (obtained by restricting the transition matrices assigned to the arcs), although a phylogenetic invariant has been described for this general model ([15]—essentially by taking determinants of the equation in (P2)). We point out here that phylogenetic invariants are, in fact, abundant for this general model since each triple or

quadruple gives rise (via (P1) and (P2), respectively) to  $r^2$  polynomial identities.

Referring to Theorem 3.1, note that, in this setting,  $H_\delta$  would not be expected to have elements of order 2, however  $\mathcal{G}$  clearly does, which is why we did not impose the simpler restriction in Theorem 3.1 that  $\mathcal{G}$  have no elements of order 2.

• Suppose  $\mathcal{G}$  is a group, and  $\mathcal{S} \subseteq \mathcal{G}$  is a semigroup. An interesting extension of Theorem 3.1 would be to characterize when a proximity map  $\delta: X \times X \rightarrow \mathcal{S}$  is a tree proximity map with arc weights lying in  $\mathcal{S}$ .

### ACKNOWLEDGMENT

We thank Andreas Dress for several helpful comments on an earlier version of this paper.

### REFERENCES

1. A. V. Aho, Y. Sagiv, T. G. Szymanski, and J. D. Ullman, Inferring a tree from lowest common ancestors with an application to the optimization of relational expressions, *SIAM J. Comput.* **10** (1981), 405–421.
2. E. Baake, What can and what cannot be inferred from pairwise sequence comparisons? *Math. Biosci.* **154** (1998), 1–21.
3. H.-J. Bandelt and A. Dress, Reconstructing the shape of a tree from observed dissimilarity data, *Adv. in Appl. Math.* **7** (1986), 309–343.
4. H.-J. Bandelt and M. A. Steel, Symmetric matrices representable by weighted trees over a cancellative abelian monoid, *SIAM J. Discrete Math.* **8**, No. 4 (1995), 517–525.
5. S. Böcker and A. W. M. Dress, Recovering symbolically dated, rooted trees from symbolic ultrametrics, *Adv. in Math.* **138** (1998), 105–125.
6. D. Bryant and M. Steel, Extension operations on sets of leaf-labelled trees, *Adv. in Appl. Math.* **16** (1995), 425–453.
7. P. Buneman, A note on the metric property of trees, *J. Combin. Theory Ser. B* **17** (1974), 48–50.
8. J. Chang, Full reconstruction of Markov models on evolutionary tree, *Math. Biosci.* **137** (1996), 51–73.
9. S. Evans and T. Speed, Invariants of some probability models used in phylogenetic inference, *Ann. Statist.* **21**, No. 1 (1993), 355–377.
10. J. S. Farris, A. G. Kluge, and M. J. Eckardt, A numerical approach to phylogenetic systematics, *Syst. Zoology* **19** (1970), 172–189.
11. S. L. Hakimi and A. N. Patrinos, The distance matrix of a graph and its tree realization, *Quart. Appl. Math.* **30** (1972), 255–269.
12. J. A. Lake, Reconstructing evolutionary trees from DNA and protein sequences: paralogous distances, *Proc. Nat. Acad. Sci. U.S.A.* **91** (1994), 1455–1459.

13. M. P. Ng and N. C. Wormald, Reconstruction of rooted trees from subtrees, *Discrete Appl. Math.* **69** (1996), 19–31.
14. J. M. S. Simões Pereira, A note on the tree realizability of a distance matrix, *J. Combin. Theory* **6** (1969), 303–310.
15. M. A. Steel, Recovering a tree from the leaf colourations it generates under a Markov model, *Appl. Math. Lett.* **7**, No. 2 (1994), 19–24.
16. K. A. Zaretskii, Constructing trees from the set of distances between pendant vertices, *Uspekhi Mat. Nauk* **20** (1965), 90–92.