

Self-generating autocatalytic networks: structural results, algorithms, and their relevance to evolutionary processes

Daniel Huson¹, Joana C. Xavier^{2,3}, Mike Steel⁴

¹Institute for Bioinformatics and Medical Informatics, Tübingen University, Tübingen, Germany.

²Dayhoff Labs, London, UK.

³Department of Chemistry, Imperial College, London, UK.

⁴Biomathematics Research Centre, University of Canterbury, Christchurch, New Zealand.

Abstract

The concept of an autocatalytic network of reactions that can form and persist, starting from just an available food source, has been formalised by the notion of a Reflexively-Autocatalytic and Food generated (RAF) set. The theory and algorithmic results concerning RAFs have been applied to a range of settings, from metabolic questions arising at the origin of life, to ecological networks, and cognitive models in cultural evolution. In this paper, we present new structural and algorithmic results concerning RAF sets, by studying more complex modes of catalysis that allow certain reactions to require multiple catalysts (or to not require catalysis at all), and discuss the differing ways catalysis has been viewed in the literature. We then focus on the structure and analysis of minimal RAFs, and derive structural results and polynomial-time algorithms, with applications to metabolic network data described briefly.

Keywords: autocatalytic network, RAFs, algorithms, metabolic network

1 Introduction

A central property of the chemistry of living systems is that they combine two basic features: (i) the ability to survive on an ambient food source and (ii) each biochemical

reaction in the system requires only reactants and a catalyst that are provided by other reactions in the system (or are present in the food set). The notion of a self-sustaining ‘collectively autocatalytic set’ captures these basic features, and their study was pioneered by Stuart Kauffman [24, 25]. By investigating a simple binary polymer model, Kauffman showed that collectively autocatalytic sets invariably emerge once the network of polymers becomes sufficiently large.

The notion of a collectively autocatalytic set was subsequently formalized more precisely as a ‘reflexively autocatalytic and food-generated’ (RAF) set (defined shortly). RAF sets (RAFTs) are related to, but somewhat different from Robert Rosen’s (M, R) systems (a partial connection between the two was described in [22]). RAF theory can also be investigated within the framework of Chemical Organisation Theory (COT) [4]; for example, certain types of RAFTs correspond to chemical organisations, as described in [19] (see also [40] Section 4).

RAF algorithms have also been used in the analysis of simple autocatalytic networks of polymers in laboratory studies, either from RNA molecules [16] or from peptides [15], and have been discussed further in modelling the origin of life (see e.g., [6, 44]). More recently, RAFTs have also played a pivotal role in modelling self-reproduction and self-organisation before the emergence of a genetic code in the polymer world. When modelling this early stage of chemical evolution [46, 47], RAFTs provide a framework for studying the organisation of small molecules in complex chemical networks that would then lead to polymers as RNA and protein. A general framework to model complex catalysis is required here: multiple small-molecules can be involved in catalysis of the same reaction, whereas some reactions can occur spontaneously. There are many examples of this in biochemistry; for example, the enzyme L-threonine dehydrogenase, important in amino acid metabolism requires an organic cofactor, namely NAD, and a metal [23]. Moreover, in the data set analysed in [46] 1052 of the 5994 biochemical reactions involved catalysts combinations of two or more molecules to be present.

Two important features of the RAF approach are the degree of generality RAFTs allow, and the fact that very large systems can be analysed precisely by fast algorithms. The generality of RAF theory means that a ‘reaction’ need not refer specifically to a chemical reaction, but to any process in which certain items are combined and transformed into new items, and where similar items facilitate (or catalyse) the process without being used up in the process. This has led to the application of RAF theory to processes beyond biochemistry, such as cognitive modelling in cultural evolution [7, 8], ecology [9, 10], and economics [10]. This generality is not unique to RAFTs; for example, Chemical Organisation Theory has been applied to diverse settings including sociology [5], ecology [11], cybernetics [13] and modelling of worldviews [45]. Petri Nets have also been applied to biochemical modelling, including self-reproduction [37] and to other non-biochemical settings (e.g. [27]). In addition, a number of other recent structural approaches to autocatalytic networks have been applied in the origin-of-life setting [2, 35].

In this paper, we describe further extensions and applications of RAF theory. We describe an extension of the RAF approach that provides a unified handling of complex

catalysis, leading to new mathematical results [Sections 2.2, 3]. We then focus on the structure and algorithmic properties of minimal RAFs in [Section 4.1].

2 Catalytic reaction systems

2.1 Reaction Systems

A *reaction system* is a pair (X, R) consisting of a finite nonempty set X of *elements* (e.g., molecule types) and a finite set R of *reactions*. Here a *reaction* $r \in R$ refers to an ordered pair (A, B) where A and B are multisets of elements from X . We will write $r : a_1 + \dots + a_k \rightarrow b_1 + \dots + b_l$ to denote the reaction that has reactants $\{a_1, \dots, a_k\}$, and products $\{b_1, \dots, b_l\}$ ¹. We let $\rho(r)$ denote the set corresponding to A (i.e., ignoring multiplicities), and $\pi(r)$ denote the set corresponding to B (ignoring multiplicities); it is assumed implicitly that $\rho(r), \pi(r) \neq \emptyset$. For a subset R' of R , it is convenient to let $\pi(R') = \bigcup_{r \in R'} \pi(r)$ denote the set of the products of the reactions in R' .

Next, consider a reaction system (X, R) together with a particular subset F of X . The set F can be interpreted as a set of elements that are freely available to the system; accordingly, F is referred to as a *food set*. A subset R' is *F-generated* if the reactions in R' can be placed in some linear order, say r_1, r_2, \dots, r_k , such that the following property holds: for $\rho(r_1) \subseteq F$ and for all values of j between 2 and k , we have $\rho(r_j) \subseteq F \cup \pi(\{r_1, \dots, r_{j-1}\})$. In other words, the reactions in R' are F-generated if they can proceed in some order so that the reactant(s) of each reaction are available by the time they are first required. We call such an ordered sequence of R' an *admissible ordering*. Since there are $k!$ ways to order k reactions, it may not be immediately obvious that the F-generated condition can be verified in polynomial time; however, there is a simple way to do so, as we now describe.

We first recall some further terminology. Given a subset R' of reactions R , a subset W of X is said to be *R'-closed* precisely when each reaction $r \in R'$ that has all its reactant(s) in W also has all its the product(s) in W (i.e. $r \in R', \rho(r) \subseteq W \Rightarrow \pi(r) \subseteq W$). The union of two closed sets need not be closed; nevertheless, given any nonempty subset W_0 of X there is a unique minimal R' -closed set containing W_0 , denoted $\text{cl}_{R'}(W_0)$. This can be computed in polynomial time in the size of the system by constructing a nested increasing sequence of subsets of the elements $W_0 \subset W_1, \dots \subset W_k$ where:

$$W_{i+1} = W_i \cup \{x \in X : \exists r \in R' : \rho(r) \subseteq W_i, x \in \pi(r)\},$$

for $i \geq 0$, and by letting W_k denote the terminal set in this sequence (i.e. k is the first value of i for which $W_i = W_{i+1}$).

2.2 Catalytic reaction systems (allowing complex catalysis)

A *catalytic reaction system* (CRS) is a reaction system with a food set (X, R, F) together with a subset χ of $2^X \times R$. Thus χ is a collection of pairs (U, r) where $U \subseteq X$ and $r \in R$. For $(U, r) \in \chi$ we refer to U as a *catalyst set* for r . This is a generalisation

¹In applications, a bidirectional reaction is generally regarded as a pair of reactions (forward and backward) with the same catalysis assignment.

of earlier treatments in which χ consisted of a subset of $X \times R$ (i.e. simple catalysis by single elements). Our extension here to this more general catalysis framework allows for complex (i.e. conjunctive) catalysis rules, where catalysts of a reaction may require the presence of two or more elements of X (e.g., cofactors of enzymes). However, the treatment of complex catalysis in [41] required the introduction of fictitious new reactions and elements to the original CRS. Here, our more direct approach allows both simple and complex catalysis rules that require no additional reactions or elements to be introduced. It also permits the further option that particular uncatalysed reactions can appear in an autocatalytic system (since the definition of χ allows $(\emptyset, r) \in \chi$), thereby addressing a recent concern discussed in Section 2.5.

We denote a CRS by writing $\mathcal{Q} = (X, R, \chi, F)$, and we let $|\mathcal{Q}| = |X| + |R| + |\chi|$ denote the *size* of \mathcal{Q} and we write

$$r : a_1 + \cdots + a_k[U_1, \dots, U_m] \rightarrow b_1 + \cdots + b_l$$

to denote the reaction that has reactants $\{a_1, \dots, a_k\}$, products $\{b_1, \dots, b_l\}$ and catalyst sets $\{U_1, \dots, U_m\}$, where $(U_i, r) \in \chi$ for all i . When U_i is a singleton set (say $\{c_i\}$), we will often write c_i in place of $\{c_i\}$ in our example systems.

2.3 RAFs

Given a CRS $\mathcal{Q} = (X, R, \chi, F)$, a subset R' of R is said to be *Reflexively-Autocatalytic and Food generated* (more briefly a *RAF*) if R' is nonempty and if for each $r \in R'$, the reactants of r and at least one catalyst set U for r (as specified by χ) is a subset of $\text{cl}_{R'}(F)$.

An equivalent definition for a nonempty set R' to be a RAF for \mathcal{Q} is that R' is F-generated, and each reaction $r \in R'$ has a catalyst set U that is a subset of $F \cup \pi(R')$. A further equivalent definition is the following:

R' can be ordered r_1, r_2, \dots, r_k so that for each $i \geq 1$, the reactants of r_i are present in X_i , where $X_1 = F$ and $X_i = F \cup \pi(\{r_1, \dots, r_{i-1}\})$, and at least one catalyst set U of r_i is a subset of X_k .

If a CRS \mathcal{Q} has a RAF, then it has a unique maximal RAF (which is the union of all the RAFs for \mathcal{Q}), which is denoted $\text{maxRAF}(\mathcal{Q})$. A RAF R for \mathcal{Q} is said to be an *irreducible* RAF (more briefly an *iRAF*) if $R \setminus \{r\}$ is not a RAF for \mathcal{Q} and contains no RAF for \mathcal{Q} .

A stronger notion than a RAF is a *CAF* (constructively-autocatalytic and F-generated set) where the third equivalent definition of a RAF (above) is strengthened to ‘and at least catalyst set U of r_i is a subset of X_i ’ (rather than ‘of X_k ’). In other words, R' is a CAF if it has an admissible ordering in which at least one catalyst set has each of its elements already present in the food set or produced by an earlier reaction in the ordering. Every CAF is also a RAF, but the converse containment does not hold. Although RAFs and CAFs appear to be very similar concepts, they exhibit quite different properties. For example, if a CRS \mathcal{Q} has a CAF, then this CAF must contain a reaction r for which all the reactants of r and at least one catalyst of r lie in F , in which case $\{r\}$ is itself a CAF of size 1. By contrast, a large RAF need not

contain any ‘small’ RAF within it. Moreover, theoretical and simulation studies on polymer systems reveal that the level of catalysis required for a CAF to be present is exponentially higher than that required for a RAF [32], and in real biochemical systems that have been studied (e.g., [38, 46]), the maxRAF is generally not a CAF. Thus, in this paper, we focus on the more general notion of a RAF.

2.4 Examples

We now describe three examples to illustrate the concepts above. The first example (from [16]) illustrates the concept of a RAF in the simpler setting where catalysis involves only singleton elements, the second example illustrates complex catalysis and the third example is from an experimental system.

Consider the following CRS where $F = \{f_1, f_2, f_3, f_4\}$, $X = F \cup \{p_1, \dots, p_6\}$, and $R' = \{r_1, \dots, r_6\}$ indicated by squares in Fig. 1. In this figure, reactants and product pathways are indicated by solids arrows, and catalysis is indicated by dashed arrows. The maxRAF for this system consists of the four reactions (r_1 – r_4), and there is one iRAF for this CRS, namely $\{r_1, r_2\}$.




Fig. 1 A simple example of a CRS with food set $\{f_1, f_2, f_3, f_4\}$ six reactions (r_1 – r_6) and with catalysis indicated via dashed arrows (adapted from [16]). The maxRAF consists of the four reactions within the blue border.

Next consider the following CRS where $X = \{a, b, c, d, e, g\}$, $F = \{a, b\}$ and $R = \{r_1, r_2, r_3, r_4\}$ and χ are as follows:

$$\begin{aligned} r_1 : a + a \ [\{c, d\}, e] &\rightarrow c, & r_2 : b + c \ [\emptyset] &\rightarrow d, \\ r_3 : b + b \ [] &\rightarrow e, & r_4 : a + e \ [a] &\rightarrow b, \\ r_5 : c + d \ [d] &\rightarrow g + g. \end{aligned}$$

Note the subtle difference between the catalyst set in r_2 and r_3 . In r_2 we have $\chi = \{\emptyset\}$ so $\emptyset \in \chi$ which means r_2 does not require a catalyst (either present in F or as a

product of another reaction), whereas for r_3 , $\chi = \emptyset$ so no catalyst set (U) exists for r_2 . This system has $\{r_1, r_2, r_5\}$ as its maxRAF, and $\{r_1, r_2\}$ as its unique iRAF. No CAF is present in this CRS.

A third example of a RAF arising in an experimental system (involving simple rather than complex catalysis) is provided in Fig. 2.




Fig. 2 *Left:* An experimental RNA system described in [43] involving 16 reactions, and 18 molecule types (including a food set of size 2, which form the reactants of each reaction). The reactions are denoted by squares, elements (in this case, RNA replicators) are denoted by circles, reaction pathways are indicated by solid arrows, and catalysis is indicated by dotted arrows. This system forms a RAF [16]. *Right:* A subset of seven of the reactions from the full set (with the two food elements combined into a single element f), which forms a RAF. This RAF, analysed in [16] contains within it 67 other subsets that form RAFs including four irreducible RAFs (iRAFs). This RAF is not a CAF (nor does it contain one). Figures produced by *CatlyNet* [21].

2.5 The interpretation of catalysis and autocatalysis in RAFs

Note that if $(\emptyset, r) \in \chi$, then r does not require any element of X for its catalysis, and so RAFs under our more general definition can contain uncatalysed reactions in settings where this is appropriate. This is particularly relevant, as a number of papers (e.g., [1, 14, 28, 26]) have pointed to the restrictive nature of RAFs in requiring that all reactions in the RAF must be catalysed. For example, the authors of [1] stated:

“Here, the assumption that all reactions are catalysed appears very unrealistic. Unfortunately, the algorithms for recognizing RAFs [...] do not seem to generalise to arbitrary networks composed of non-catalysed reactions.”

and in [14] the author states:

“The requirement that every reaction be catalyzed by another molecule seems too strong when we are dealing with simple metabolisms in the earliest forms of life. In modern organisms, almost every reaction of small-molecule metabolites is catalyzed by enzymes. However, prior to the existence of RNA and proteins, there were no enzymes, so we need a

theory that deals with the reactions of small molecules without insisting that the reactions be catalyzed.”

In fact, in previous applications of RAF theory to the origins of metabolism [46, 47] a fictional catalyst ‘Spontaneous’ was assigned to reactions known to occur uncatalysed. This catalyst was added to the food set in all simulations. Also, prior to the advent of genetic coding and enzymes, catalysis must have existed in the small-molecule world as well. Small molecules are increasingly being shown to catalyse multiple reactions in the absence of enzymes [3, 33]. However, the use of a fictional catalyst ‘Spontaneous’ is less direct than our approach introduced here (see Section 2.2) where we generalise the notion of catalysis by describing it via a subset χ of $2^X \times R$ (thereby allowing pairs of the form (\emptyset, r) , and so a ‘Spontaneous’ catalyst is no longer required).

RAF theory also treats ‘catalysis’ in a general way and this allows for efficient graph-theoretic algorithms, which apply independently of any detailed kinetic (or even stoichiometric) considerations. This generality also allows for applications in a variety of areas outside chemistry. Essentially, we regard a catalyst as any element that facilitates, speeds-up, or synchronises a reaction without taking part in the reaction itself (as a reactant).

For example, in economic applications, a factory facilitates (i.e. catalyses) the production of items from the incoming raw materials, but is not itself consumed by that process. In cognitive modelling [7], a reaction that combines ideas to form a new idea could be enhanced (i.e. catalysed) by a need, thought, memory, or stimuli. In ecology [9], a catalyst is a species that enables some other interaction in an ecological network. In biochemistry, it can also be helpful to treat catalysis in a quite general way; for instance, the formation of a lipid boundary to encompass a primitive metabolism can be viewed as a sequence of reactions which, once complete, forms a structural element (a complete lipid membrane) that catalyses all the reactions within the newly formed protocell (since the system within it no longer disperses) [17].

Moreover, in biochemistry, food elements can be direct catalysts for multiple reactions, as is the case with universally-essential metal ions. The generality of RAF theory allows for this, in contrast to some other models of network autocatalysis.

It is tempting to treat a catalyst x of a reaction r as simply an additional reactant and product of r (i.e., adding x to both sides of the reaction, as is sometimes used in chemical notation), thereby reducing the notion of a RAF to simply an F-generated subset of the resulting (modified) reaction system. In other words, the catalysed reaction $x + y[c] \rightarrow z$ (where c is a catalyst) might be viewed as $x + y + c \rightarrow z + c$. However, this misses an important distinction. Namely, in RAF theory, it is assumed that a reaction r may proceed (at a slow rate initially) provided that all its reactants are available even if no catalyst is initially available; however r will subsequently be part of a RAF set, provided that a catalyst of r is (eventually) produced by at least one other reaction in the RAF. Indeed, it is entirely possible for a CRS to contain a RAF yet the modified reaction system (adding catalysts to both sides of a reaction) might have no F-generated subset (indeed, this has been observed in real biochemical networks [16, 38, 46]). Misunderstandings concerning the related notion of ‘autocatalysis’ have been discussed in [18] and [36].

3 Mathematical aspects of RAFs

We now recall the concept of the maxRAF operator φ (from [42], Section 3). For any subset R' of R , let $\mathcal{Q}|R'$ be the CRS (X, R', χ', F) where χ' is the restriction of χ to $2^X \times R'$, and let $\varphi : 2^R \rightarrow 2^R$ be the function defined as follows:

$$\varphi(R') = \begin{cases} \text{maxRAF}(\mathcal{Q}|R'), & \text{if } \mathcal{Q}|R' \text{ has a RAF;} \\ \emptyset, & \text{otherwise.} \end{cases} \quad (1)$$

The function $\varphi(R')$ can be determined in a computationally efficient way, as follows:

$$\varphi(R') = \bigcap_{i \geq 1} R_i,$$

where $R_1 = R'$ and for $i > 1$,

$$R_i = \{r \in R_{i-1} : \rho(r) \subseteq \text{cl}_{R_{i-1}}(F) \text{ and } (U, r) \in \chi \text{ for some } U \subseteq \text{cl}_{R_{i-1}}(F)\}. \quad (2)$$

Notice that the sets R_i defined in Eqn. (2) form a decreasing nested sequence, and so $\varphi(R')$ is precisely the set R_j for the first value of $j \geq 1$ for which $R_{j+1} = R_j$. In particular, a nonempty subset R' of R is a RAF if and only if $\varphi(R') = R'$. Moreover, $\varphi(R)$ is precisely the maxRAF of \mathcal{Q} (if it exists), or is the empty set otherwise.

Example: To illustrate this algorithm, consider the second example described in Section 2.4. For $R' = R = \{r_1, \dots, r_5\}$ we have $R_1 = R$, and so $\text{cl}_{R_1}(F) = \{a, b, c, d, e, g\}$, and all but one of the reactions (namely, r_3) satisfy the property that $(U, r) \in \chi$ for some $U \subseteq \text{cl}_{R_1}(F)$. Thus, (by Eqn. (2)), $R_2 = \{r_1, r_2, r_4, r_5\}$. Next, $R_3 = \{r_1, r_2, r_5\}$ since one of the reactants of $r_4 \in R_2$ (namely e) is not present in $\text{cl}_{R_2}(F)$. Finally, $R_4 = R_3$, and so $\varphi(R) = R_3$ forms the maxRAF for \mathcal{Q} .

A number of interesting algebraic (semigroup) properties of the map φ have been explored recently in [29] (see also [30], which considers a more general notion than a RAF, which corresponds to what have been called ‘pseudo-RAFTs’ in the RAF literature).

The map φ is also an example of the more general notion of an interior operator (on 2^R), and several basic results in RAF theory can be derived by this property alone. To explain this further, an arbitrary function $\psi : 2^Y \rightarrow 2^Y$ is said to be an *interior operator*² if it satisfies the following three properties (nesting, monotonicity, and idempotence) for all subsets A, A' of Y :

- (I₁): $\psi(A) \subseteq A$,
- (I₂): $A \subseteq A' \Rightarrow \psi(A) \subseteq \psi(A')$, and
- (I₃): $\psi(\psi(A)) = \psi(A)$.

²The term ‘interior operator’ comes from topology, since the function that assigns to any subspace S of a topological space the topological interior of S satisfies the three properties described.

Lemma 1. *Given any CRS $\mathcal{Q} = (X, R, \chi, F)$ the maxRAF function $\varphi : 2^R \rightarrow 2^R$ is an interior operator.*

Many results of RAF theory (including some we discuss later) depend mainly on the interior operator property of φ . A natural question is whether *any* interior operator on a finite set can be represented as the maxRAF operator of a set of reactions associated with the elements of the set. It was recently shown that if Y is a finite set of size at least 12, then there exist interior operator on Y that cannot be represented as a maxRAF operator of a suitably chosen CRS (X, R_Y, χ, F) where R_Y is a set of reactions bijectively associated with Y and where $\chi \subseteq X \times R_Y$ (i.e. the original CRS setting which does not allow complex catalysis rules) [39]. Thus some generic properties of RAFs in this setting cannot be established by using the interior operator properties alone. However, if one allows complex catalysis rules, we have the following contrasting result (further details and a proof are provided in the Appendix).

Proposition 1. *For any finite set Y , any interior operator ψ on 2^Y can be represented as the maxRAF operator φ for a suitably chosen CRS $\mathcal{Q} = (X, R_Y, \chi, F)$, where R_Y is bijectively associated with Y and $\chi \subseteq 2^X \times R_Y$ permits complex catalysis rules. Moreover, R_Y can be chosen so that each of its reactions has the same single reactant f (from the food set) and a single associated product.*

Proposition 1 has an interesting consequence for the structure of RAFs in any CRS. Specifically, it implies that it is possible to simplify the reactions and the food set, while preserving the number, sizes, and containment structure of the RAFs (albeit at the price of possibly increasing substantially the number and complexity of catalysts for the reactions). More precisely, we have the following result (obtained from Proposition 1 by taking ψ to be the maxRAF operator of \mathcal{Q}).

Corollary 1. *For any CRS $\mathcal{Q} = (X, R, \chi, F)$, there is a matching CRS $\mathcal{Q}' = (X', R', \chi', F')$, where (i) each of the reactions in R' has a single reactant f which comprises the food set F , and a single distinct product and (ii) there is a canonical bijective correspondence between R and R' that induces a bijection between the RAFs of \mathcal{Q} and the RAFs of \mathcal{Q}' .*

A worked example to illustrate this result is provided in the Appendix. We return to interior operators in Section 4.2.


3.1 Strictly autocatalytic RAFs

If a CRS \mathcal{Q} has the property that each reaction is catalysed by at least one element of the food set, then the RAF sets for \mathcal{Q} coincide precisely with the F -generated sets. More generally, for any CRS, it is possible for a RAF set to have the property that some (or all) of its reactions are catalysed by elements of the food set, which renders the notion of ‘autocatalytic’ less applicable (though not entirely, since each reaction in the RAF might still also be catalysed by at least one product of the other reactions in

the RAF). To formalise this notion further, we introduce a new definition and describe a result that characterises the condition which captures a more focused notion of an ‘autocatalytic’ RAF.

Given a CRS $\mathcal{Q} = (X, R, \chi, F)$, we say that a subset R' of R is *strictly autocatalytic* if each reaction in R' has at least one catalyst that involves one or more products of reactions of R' and is not a subset of the food set. We say R' is a strictly autocatalytic RAF for \mathcal{Q} if R' is a RAF and is strictly autocatalytic.

Example: Consider the following catalytic reaction system:



where $X = \{a, b, c, d, e\}$ and $F = \{a, b, c\}$. Then, $\{r_2\}$, $\{r_1, r_2\}$, $\{r_2, r_3\}$ and $\{r_1, r_2, r_3\}$ are RAFs, however $\{r_1, r_2\}$ is the only strictly autocatalytic RAF. \square

Let $\hat{R}_i, i \geq 1$ be the nested decreasing sequence of subsets of R defined as follows: $\hat{R}_1 = R$, and for $i > 1$, consider the following modification of Eqn. (2):

$$\hat{R}_i = \{r \in \hat{R}_{i-1} : \rho(r) \subseteq \text{cl}_{\hat{R}_{i-1}}(F) \text{ and } (U, r) \in \chi \text{ for some } U \subseteq \text{cl}_{\hat{R}_{i-1}}(F) \text{ and } U \not\subseteq F\}. \quad (3)$$

The following result shows that there is a polynomial-time algorithm (in $|\mathcal{Q}|$) to determine whether or not \mathcal{Q} has a strictly autocatalytic RAF and, if so, to construct a maximal one. The proof is provided in the Appendix.

Proposition 2. *Let $\mathcal{Q} = (X, R, \chi, F)$ be a CRS. Then \mathcal{Q} has a strictly autocatalytic RAF if and only if $\bigcap_{i \geq 1} \hat{R}_i \neq \emptyset$, in which case, $\bigcap_{i \geq 1} \hat{R}_i$ is the unique maximal strictly autocatalytic RAF for \mathcal{Q} .*

Application: We investigated the large primitive metabolic data-set used in [46] based on 6039 reactions and a food set of size 68, which resulted in a maxRAF of size 1357. Applying the strictly autocatalytic RAF algorithm determined that no such RAF exists. This, in turn, implies that in any RAF for this system, one or more elements in the food set play a pivotal role in catalysing some reaction(s). This is consistent with the aforementioned essentiality of metals in biocatalysis of central metabolism. At the same time, this result reveals that there are unknown prebiotic routes (or unknown non-enzymatic catalysts) to the production of organic cofactors which must be investigated in the laboratory. For example, there are no routes in this network for the production of organic cofactors as NAD [47] and the investigation of the prebiotic route for its synthesis is subject of active investigation [12, 34].

4 Irreducible (minimal) RAFs

Recall that an irreducible RAF (iRAF) is a RAF that contains no RAF as a proper subset. Such RAFs are of particular interest, as they represent the minimal possible autocatalytic networks within a CRS. A smallest sized RAF in any CRS is necessarily an iRAF, however, iRAFs can be of different sizes. Moreover, even without complex catalysis rules, a CRS \mathcal{Q} can have exponentially many iRAFs, and although finding one iRAF is easy, finding a smallest one turns out to be NP-hard [40].

4.1 Identifying all iRAFs in a CRS

Let $\mathcal{Q} = (X, R, \chi, F)$ be a fixed CRS that contains a RAF. Recall that a RAF R for \mathcal{Q} is said to be an iRAF (irreducible RAF) if $R \setminus \{r\}$ is not a RAF for \mathcal{Q} and contains no RAF for \mathcal{Q} . Every RAF of a CRS contains an iRAF, and there is a simple (polynomial-time in $|\mathcal{Q}|$) algorithm for finding one or more iRAFs within any given RAF; however, the number of iRAFs for \mathcal{Q} can grow exponentially with $|\mathcal{Q}|$ (Theorem 1 of [20]). Moreover, determining the size of a *smallest* iRAF is known to be NP-hard [40]. Nevertheless, there is a simple (and polynomial-time) algorithm to determine whether \mathcal{Q} has just one iRAF; we simply ask whether the set

$$\text{Core}(\mathcal{Q}) := \{r \in R : \varphi(R \setminus \{r\}) = \emptyset\}$$

is a RAF. Then it can be shown that this set is a RAF if and only if \mathcal{Q} has a unique iRAF, and in that case, $\text{Core}(\mathcal{Q})$ is the unique iRAF for \mathcal{Q} . The result for CRS systems without complex catalysis was established in see [42] Theorem 4.1, however since the proof involved only interior operator properties for φ it applies to the more general setting here (by Lemma 1).




Moreover, it can easily be seen that there is a simple way to test whether or not an iRAF R_1 is the only iRAF for \mathcal{Q} ; this holds provided that:

$$\varphi(\varphi(R) \setminus \{r\}) = \emptyset, \text{ for all } r \in R_1.$$

This is computationally less-intensive than computing $\text{Core}(\mathcal{Q})$, since it involves searching over the reactions that are just in R_1 rather than all of R .

In a similar way, it can be shown that if \mathcal{Q} has (at least) two iRAFs, R_1 and R_2 then these are the only iRAFs for \mathcal{Q} if and only if for all $r \in R_1 \setminus R_2$, we have $\varphi(\varphi(R) \setminus \{r\}) = R_2$, and for all $r \in R_2 \setminus R_1$, we have $\varphi(\varphi(R) \setminus \{r\}) = R_1$.

A direct extension of this last result to three or more iRAFs is problematic, since although two iRAFs cannot be nested (i.e. neither can be a subset of the other) in the case of three iRAFs, it is possible for each one to be a subset of the union of the two others. An example is provided by the CRS $\mathcal{Q} = (X, R, \chi, F)$, where $F = \{f\}$, $R = \{r_1, r_2, r_3\}$ and



This system has three iRAFs, namely, $\{r_1, r_2\}$, $\{r_1, r_3\}$, and $\{r_2, r_3\}$, each of which contained in the union of the other two.

However, the following result shows that testing a small number (say, k) of iRAFs is feasible in polynomial time in $|\mathcal{Q}|$ (provided that k is fixed). The result is a slight strengthening of Theorem 2 of [40]; its significance is that it allows one to determine whether a given set of k iRAFs is the set of *all* iRAFs for \mathcal{Q} in polynomial time in $|\mathcal{Q}|$, provided that k is fixed (however, the algorithm is exponential in k). The proof of the following result is given in the Appendix.

Proposition 3. *A given collection R_1, \dots, R_k of iRAFs for \mathcal{Q} is the set of all iRAFs for \mathcal{Q} if and only if the following condition holds: For all choices (r_1, \dots, r_k) where $r_i \in R_i$, we have $\varphi(\varphi(R) \setminus \{r_1, \dots, r_k\}) = \emptyset$.*

Proposition 3 assumes that the iRAFs are given in advance (e.g., found by heuristic search); however, the approach can be extended to identify the complete set of iRAFs of \mathcal{Q} even if they are not given in advance.

Thus, when the number of iRAF is small (say, less than some value k), the following algorithm is polynomial-time in $|\mathcal{Q}|$ (but is exponential in k). The algorithm proceeds as follows and when it terminates, it produces the complete list of iRAFs of \mathcal{Q} .

- Given $\mathcal{Q} = (X, R, \chi, F)$, determine the maxRAF; providing that it exists, compute an iRAF, denoted R_1 .
- For $i \geq 1$, construct a sequence of iRAFs starting with R_1 , as follows.
 - (i) For each integer j with $1 \leq j \leq i$ and each choice of i reactions $\{r_1, r_2, \dots, r_i\}$ where $r_j \in R_j$ for each j , compute $\varphi(\varphi(R) \setminus \{r_1, \dots, r_i\})$. If this set is the empty set for all such choices of $\{r_1, \dots, r_i\}$, then R_1, \dots, R_i is the complete set of iRAFs for \mathcal{Q} , so the algorithm terminates.
 - (ii) Otherwise, if $\varphi(\varphi(R) \setminus \{r_1, \dots, r_i\}) \neq \emptyset$ for some choice of $\{r_1, \dots, r_i\}$, then compute an iRAF of $\varphi(\varphi(R) \setminus \{r_1, \dots, r_i\})$ and set R_{i+1} equal to this iRAF, and proceed to step (iii).
 - (iii) Repeat step (i) and, if necessary, step (ii).

4.2 Finding an iRAF that contains a given reaction

Given any CRS $\mathcal{Q} = (X, R, \chi, F)$, a natural question is whether a given reaction in R is present in at least one iRAF for \mathcal{Q} . We show shortly that this problem is NP-hard. First, we describe how the union of all the iRAFs present in a CRS can be described via a further interior operator. Given a CRS $\mathcal{Q} = (X, R, \chi, F)$, and a nonempty subset R' of R , let $\tilde{\varphi}(R')$ denote the union of the subsets of R' that are iRAFs for \mathcal{Q} . The proof of the following lemma is given in the Appendix, as a consequence of a more general result.

Lemma 2. *The map $\tilde{\varphi} : 2^R \rightarrow 2^R$ is an interior operator.*

Although the computation of $\varphi(R')$ is polynomial time in $|\mathcal{Q}|$ for any CRS $\mathcal{Q} = (X, R, \chi, F)$ and any subset R' of R , computing $\tilde{\varphi}(R)$ is NP-complete, even for

quite simple CRS systems, as we now state more formally (a proof is provided in the Appendix).

Proposition 4. *Given a CRS $\mathcal{Q} = (X, R, \chi, F)$ the problem of determining whether or not a given reaction $r \in R$ is present in at least one iRAF for \mathcal{Q} is NP-complete. Moreover, this holds even for systems where each reaction in R has the same single food-set reactant and where each catalyst consists of single elements (i.e. without complex catalysis).*


Next suppose that a CRS $\mathcal{Q} = (X, R, \chi, F)$ and that an element $x \in X \setminus F$ is produced by some reaction in $\max\text{RAF}(\mathcal{Q})$. A relevant question in certain applications is to find a minimal subset R' of reactions within $\max\text{RAF}(\mathcal{Q})$ that is both a RAF and produces one or more particular elements of interest (e.g., it plays a key role in a metabolic network, such as ATP or NAD, the universal currencies of energy and redox power in metabolism, respectively). In other words, R' is a RAF that produces x , and every proper subset of R' either fails to be a RAF or fails to produce all of the specified elements. It turns out that finding such a minimal set has a polynomial-time solution, as we now show (the proof is provided in the Appendix).

Proposition 5. *Suppose that one or more particular elements $x_1, \dots, x_k \in X \setminus F$ are produced by some reaction in $\max\text{RAF}(\mathcal{Q})$. Let $\mathcal{Q}^{+\{x_1, \dots, x_k\}}$ be the CRS obtained from \mathcal{Q} by replacing each catalyst $(U, r) \in \chi$ with $(U \cup \{x_1, \dots, x_k\}, r)$. Then the collection of minimal subsets of $\max\text{RAF}(\mathcal{Q})$ that are simultaneously RAFs of \mathcal{Q} and produce each of the elements x_1, \dots, x_k are precisely the iRAFs of $\mathcal{Q}^{+\{x_1, \dots, x_k\}}$.*

As an example with the 16-reaction CRS shown in Fig. 2, for each element x of $X \setminus F$ the minimal subset that is both a RAF and generates x has size either 1 or 2.

4.3 Describing a RAF in terms of a composition sequence involving iRAFs

Any RAF contains the union of its iRAFs; however, it may be strictly larger than this union. A simple example is provided by the simple catalytic reaction system:



where $F = \{f\}$ and $X = \{f, y, z\}$. This system is a RAF but its only iRAF is $\{r_1\}$. In general, a RAF R' is the union of its iRAFs if and only if each reaction in R' is contained in an iRAF.

Nevertheless, any RAF R' in a CRS $\mathcal{Q} = (X, R, \chi, F)$ can be described by a sequence of iRAFs and associated catalytic reaction systems. This is loosely analogous to the description of finite groups in abstract algebra via a ‘composition series’, in which a group is reduced to the trivial group via quotients that are simple groups.

Here the analogue of a group (respectively, a simple group) is a subset of reactions (respectively, an iRAF), and the analogue of a quotient group is the complement of a iRAF in a set of reactions. However, this analogy is only suggestive; for a finite group the set of associated simple groups is uniquely determined by the group, however, we do not expect the same uniqueness to hold concerning the set of associated iRAFs in our decomposition.

To describe this in our setting, suppose we have a CRS, $\mathcal{Q} = (X, R, \chi, F)$ and any nonempty subset $R' \subseteq R$. A *composition sequence* for R' is a nested decreasing sequence of subsets of R' , R_1, \dots , with $R_1 = R'$ and with $R_{i+1} = R_i \setminus \hat{R}_i$ where \hat{R}_i is an iRAF of $\mathcal{Q}_i = (X, R_i, \chi, F \cup \pi(\hat{R}_{i-1}))$ for each $i \geq 1$ for which \mathcal{Q}_i has a (nonempty) RAF. Since the sets R_i form a nested decreasing sequence of sets, we refer to the final distinct set (i.e. $\bigcap_{i \geq 1} R_i$) as the *terminal set* of the sequence.

It is clear that every subset R' of R has a composition sequence, and that this can be constructed in polynomial time in $|\mathcal{Q}|$. Moreover, by definition, the iRAF sets \hat{R}_i are disjoint. In the simple 2-reaction example above, the unique composition sequence for $R' = \{r_1, r_2\}$ is: $\{r_1, r_2\}, \{r_2\}, \emptyset$. The proof of the following result is provided in the Appendix.

Proposition 6. *Let $\mathcal{Q} = (X, R, \chi, F)$ be a CRS, let R' be a nonempty subset of R , and let R_1, R_2, \dots be a composition sequence for R' . The following are equivalent:*

- (i) R' is a RAF for \mathcal{Q} ;
- (ii) R_1, R_2, \dots has terminal set \emptyset .
- (iii) The associated iRAF sets $(\hat{R}_i, i \geq 1)$ partition R' .

Note that the iRAF sets \hat{R}_i in (iii) are not, in general, iRAFs of the original CRS \mathcal{Q} , and since the sets \hat{R}_i are (by definition) disjoint, the partitioning condition in (iii) is equivalent to the condition that the union of the sets \hat{R}_i is equal to R' .

5 Concluding comments

Autocatalytic networks have provided a formal tool to investigate processes in early biochemistry and certain other settings that involve the formation and evolution of complex structures. The discrete nature of the model provides a way to develop and implement efficient algorithms that can be applied to large data sets, and to elucidate their structural properties, such as their building blocks in terms of iRAFs or the ordering of reactions.

In this paper, we have developed further techniques that open the door to more detailed investigations into the structural properties of RAFs and iRAFs, their extensions to more general catalytic scenarios, and the investigation of the constraints on the order of reactions and the appearance of particular elements of interest. Several of the techniques described here have been implemented in the open-source software package *CatlyNet* [21], which we plan to apply to investigate further aspects of early metabolism and related evolutionary questions.

6 Acknowledgements

We thank the Royal Society Te Apārangi (New Zealand) for funding under the Catalyst Leader programme (agreement No. ILF-UOC1901). We also thank the two anonymous reviewers for numerous helpful comments and suggestions on an earlier version of this manuscript.

References

- [1] J. L. ANDERSEN, C. FLAMM, D. MERKLE, AND P. F. STADLER, *Defining autocatalysis in chemical reaction networks*. Arxiv: 2107.03086, 2021.
- [2] A. BLOKHUIS, D. LACOSTE, AND P. NGHE, *Universal motifs and the diversity of autocatalytic systems*, Proceedings of the National Academy of Sciences, 117 (2020), pp. 25230–25236, <https://doi.org/10.1073/pnas.2013527117>.
- [3] Q. DHERBASSY, R. J. MAYER, K. B. MUCHOWSKA, AND J. MORAN, *Metal-pyridoxal cooperativity in nonenzymatic transamination*, J. Am. Chem. Soc., 145 (2023), pp. 13357–13370.
- [4] P. DITTRICH AND P. SPERONI DI FENIZIO, *Chemical organization theory*, Bulletin of Mathematical Biology, 69 (2007), pp. 1199–1231.
- [5] P. DITTRICH AND L. WINTER, *Chemical organizations in a toy model of the political system*, Advances in Complex Systems, 11 (2008), p. 10:14766.
- [6] A. FILISSETTI, M. VILLANI, C. DAMIANI, A. GRAUDENZI, A. ROLI, W. HORDIJK, AND R. SERRA, *On RAF sets and autocatalytic cycles in random reaction networks*, in Advances in Artificial Life and Evolutionary Computation, C. Pizzuti and G. Spezzano, eds., Cham, 2014, Springer International Publishing, pp. 113–126.
- [7] L. GABORA AND M. STEEL, *Autocatalytic networks in cognition and the origin of culture*, J. Theor. Biol., 431 (2017), pp. 87–95.
- [8] L. GABORA AND M. STEEL, *A model of the transition to behavioral and cognitive modernity using reflexively autocatalytic networks*, J. R. Soc. Interface, 17 (no. 20200545) (2020), pp. 1–14, <https://doi.org/10.1098/rsif.2020.0545>.
- [9] R. C. GATTI, B. FATH, W. HORDIJK, S. KAUFFMAN, AND R. ULANOWICZ, *Niche emergence as an autocatalytic process in the evolution of ecosystems*, Journal of Theoretical Biology, 454 (2018), pp. 110–117.
- [10] R. C. GATTI, R. KOPPL, B. D. FATH, S. KAUFFMAN, W. HORDIJK, AND R. E. ULAMNOWICZ, *On the emergence of ecological and economic niches*, Journal of Bioeconomics, 22 (2020), pp. 99–127.
- [11] C. GAUCHEREL AND F. POMMEREAU, *Using discrete systems to exhaustively characterize the dynamics of an integrated ecosystem*, Methods in Ecology and Evolution, 10 (2019), pp. 1615–1627, <https://doi.org/10.1111/2041-210X.13242>.
- [12] J. E. GOLDFORD, H. B. SMITH, L. M. LONGO, B. A. WING, AND S. E. MCGLYNN, *Continuity between ancient geochemistry and modern metabolism enabled by non-autocatalytic purine biosynthesis*, bioRxiv, (2022), <https://doi.org/10.1101/2022.10.07.511356>, <https://www.biorxiv.org/content/early/2022/10/07/2022.10.07.511356>.

- [13] F. HEYLIGHEN AND C. JOSLIN, *Cybernetics and second order cybernetics*, in Encyclopedia of Physical Science and Technology (vol. 4), Academic Press, New York, 2003, pp. 155–170, <https://doi.org/10.1016/B0-12-227410-5/00161-7>.
- [14] P. G. HIGGS, *When is a reaction network a metabolism? criteria for simple metabolisms that support growth and division of protocells*, Life, 11 (2021), p. 966, <https://doi.org/10.3390/life11090966>.
- [15] W. HORDIJK, S. SHICHOR, AND G. ASHKENASY, *The influence of modularity, seeding, and product inhibition on peptide autocatalytic network dynamics*, ChemPhysChem, 19 (2018), pp. 2437–2444.
- [16] W. HORDIJK AND M. STEEL, *A formal model of autocatalytic sets emerging in an RNA replicator system*, J. Syst. Chem., 4 (2013), pp. 1–9.
- [17] W. HORDIJK AND M. STEEL, *Autocatalytic sets and boundaries*, Journal of Systems Chemistry, 6 (2015), pp. 1–5, <https://doi.org/10.1186/s13322-014-0006-2>.
- [18] W. HORDIJK AND M. STEEL, *Chasing the tail: The emergence of autocatalytic networks*, Biosystems, 152 (2017), pp. 1–10.
- [19] W. HORDIJK, M. STEEL, AND P. DITTRICH, *Autocatalytic sets and chemical organizations: Modeling self-sustaining reaction networks at the origin of life*, New J. Phys., 20 (no. 015011) (2018), pp. 1–13.
- [20] W. HORDIJK, M. STEEL, AND S. KAUFFMAN, *The structure of autocatalytic sets: Evolvability, enablement, and emergence*, Acta Biotheor., 60 (2012), pp. 379–392.
- [21] D. H. HUSON AND M. STEEL, *Catlynet*, 2023, <https://github.com/husonlab/catlynet>.
- [22] S. JARAMILLO AND ET AL, *(M,R) systems and RAF sets: common ideas, tools and projections*, in in Proc. ALIFEXII Conf., Odense, Denmark, August 2010, 17, MIT Press, Cambridge, MA, 2010, pp. 94–100.
- [23] A. R. JOHNSON, Y. W. CHEN, AND E. E. DEKKER, *Investigation of a catalytic zinc binding site in Escherichia coli L-threonine dehydrogenase by site-directed mutagenesis of cysteine-38*, Arch Biochem Biophys., 358 (1998), pp. 211–221, <https://doi.org/10.1006/abbi.1998.0845>.
- [24] S. A. KAUFFMAN, *Autocatalytic sets of proteins*, J. Theor. Biol., 119 (1986), pp. 1–24, [https://doi.org/10.1016/S0022-5193\(86\)80047-9](https://doi.org/10.1016/S0022-5193(86)80047-9).
- [25] S. A. KAUFFMAN, *The Origins of Order*, Oxford University Press, New York, NY, USA, 1993.
- [26] D. LANCET, R. ZIDOVETZKI, AND O. MARKOVITCH, *Systems protobiology: origin of life in lipid catalytic networks*, J. Roy. Soc. Interface, 15 (no. 20180159) (2018), pp. 1–36, <https://doi.org/10.1098/rsif.2018.0159>.
- [27] X. LI, W. YU, AND F. LARA-ROSANO, *Dynamic knowledge inference and learning under adaptive fuzzy petri net framework*, IEEE Transactions on Systems, Man, and Cybernetics, Part C (Applications and Reviews), 30 (2000), pp. 442–450, <https://doi.org/10.1109/5326.897071>.
- [28] Y. LIU, *On the definition of a self-sustaining chemical reaction system and its role in heredity*, Biology Direct, 15 (2020), p. 15, <https://doi.org/10.1186/s13062-020-00269-0>.
- [29] D. LOUTCHKO, *An algebraic characterization of self-generating chemical reaction*

- networks using semigroup models*, J. Math. Biol., 86 (2023), pp. 1–36, <https://doi.org/10.1007/s00285-023-01899-4>.
- [30] D. LOUTCHKO, *Semigroup models for biochemical reaction networks*, J. Math. Biol., 86 (2023), p. 78, <https://doi.org/10.1007/s00285-023-01898-5>.
 - [31] A. LUBIW, *A note on odd/even cycles*, Discrete Applied Mathematics, 22 (1988), pp. 87–92.
 - [32] E. MOSSEL AND M. STEEL, *Random biochemical networks: the probability of self-sustaining autocatalysis*, J. Theor. Biol., 233 (2005), pp. 327–336.
 - [33] K. B. MUCHOWSKA, S. J. VARMA, AND J. MORAN, *Nonenzymatic metabolic reactions and life’s origins*, Chem. Rev., 120 (2020), pp. 7708–7744.
 - [34] K. PASCHEK, M. LEE, D. A. SEMENOV, AND T. K. HENNING, *Prebiotic vitamin b3 synthesis in carbonaceous planetesimals*, ChemPlusChem, n/a (2023), p. e202300508.
 - [35] Z. PENG, J. LINDEROTH, AND D. A. BAUM, *The hierarchical organization of autocatalytic reaction networks and its relevance to the origin of life*, PLoS Comput Biol, 18 (no. e1010498) (2022), pp. 1–28.
 - [36] Z. PENG, K. PASCHEK, AND J. C. XAVIER, *What Wilhelm Ostwald meant by “autokatalyse” and its significance to origins-of-life research*, BioEssays, 44 (no. 2200098) (2022).
 - [37] A. A. SHAROV, *Self-reproducing systems: structure, niche relations and evolution*, Biosystems, 25 (1991), pp. 237–249.
 - [38] F. L. SOUSA, W. HORDIJK, M. STEEL, AND W. F. MARTIN, *Autocatalytic sets in E. coli metabolism*, Journal of Systems Chemistry, 6 (no. 4) (2015), pp. 1–21.
 - [39] M. STEEL, *Interior operators and their relationship to autocatalytic networks*, Acta Biotheoretica, 71: 21 (2023).
 - [40] M. STEEL, W. HORDIJK, AND J. SMITH, *Minimal autocatalytic networks*, Journal of Theoretical Biology, 332 (2013), pp. 96–107.
 - [41] M. STEEL, W. HORDIJK, AND J. C. XAVIER, *Autocatalytic networks in biology: structural theory and algorithms*, J. Royal Society Interface, 16 (no. 20180808) (2019), pp. 1–10.
 - [42] M. STEEL, J. C. XAVIER, AND D. H. HUSON, *The structure of autocatalytic networks, with application to early biochemistry*, J. Royal Society Interface, 17 (n. 20200488) (2020), pp. 1–11.
 - [43] N. VAIDYA, M. L. MANAPAT, I. A. CHEN, R. XULVI-BRUNET, E. J. HAYDEN, AND N. LEHMAN, *Spontaneous network formation among cooperative RNA replicators*, Nature, 491 (2012), pp. 72–77.
 - [44] V. VASAS, C. FERNANDO, M. SANTOS, S. KAUFFMAN, AND E. SZATHMÁRY, *Evolution before genes*, Biology Direct, 7 (no. 1) (2012), pp. 1–14.
 - [45] T. VELOZ AND P. MALDONADO, *Modelling worldviews as stable metabolisms*, Entropy, 17 (2022), p. 10:14766.
 - [46] J. C. XAVIER, W. HORDIJK, S. KAUFFMAN, M. STEEL, AND W. F. MARTIN, *Autocatalytic chemical networks at the origin of metabolism*, Proc. Roy. Soc. B., 287 (no. 20192377) (2020), pp. 1–10, <https://doi.org/10.1098/rspb.2019.2377>.
 - [47] J. C. XAVIER AND S. KAUFFMAN, *Small-molecule autocatalytic networks are universal metabolic fossils*, Phil. Trans. Roy. Soc. A, 380 (no. 20210244) (2022).

7 Appendix: Mathematical details

Proof of Lemma 1: For a subset R' of R , let

$$\lambda(R') = \{r \in R' : \rho(r) \subseteq \text{cl}_{R'}(F) \text{ and } (U, r) \in \chi \text{ for some } U \subseteq \text{cl}_{R'}(F)\}.$$

It is clear that the function λ satisfies properties (I_1) and (I_2) (but not necessarily (I_3)), and since $\varphi(R') = \bigcap_{i \geq 1} H_i(R')$ where $H_1(R') = R'$ and $H_{i+1}(R') = \lambda(H_i(R'))$ for all $i \geq 1$, Property (I_3) holds by Proposition 1 of [39]. \square

Proof of Proposition 1: Let ψ be an arbitrary interior operator on 2^Y . We construct a CRS $\mathcal{Q}_\psi = (X, R_Y, \chi, F)$ that permits complex catalysis rules, as follows. Let $F = \{f\}$, let $X = \{x_y : y \in Y\} \cup \{f\}$, and let $R_Y = \{r_y : y \in Y\}$ where r_y is the reaction $f \rightarrow x_y$. It remains to describe, for each reaction $r_y \in R_Y$, its associated catalyst sets (i.e. the subsets U of X for which $(U, r_y) \in \chi$). Let

$$C(y) = \{A \subseteq Y : y \in A, \psi(A) = A\}, \quad (4)$$

the fixed sets of ψ that contain y . Then we let (U, r_y) be an element of χ if and only if $U = \{x_a : a \in A\}$ for some set $A \in C(y)$.

Let $b : Y \rightarrow R_Y$ denote the bijection $y \mapsto r_y$.

Claim 1: A nonempty set A of Y is a fixed set of ψ if and only if $b(A) = \{r_y : y \in A\}$ is a RAF for \mathcal{Q}_ψ .

To establish this claim, suppose first that A is a fixed set of ψ (i.e. $\psi(A) = A$). Since any nonempty subset of R_Y is F -generated (because each reaction in R_Y has only the single reactant f which lies in the food set), it suffices to show that each reaction r_y in $b(A)$ has an associated catalyst set U with the property that each of its elements is produced by at least one reaction in $b(A)$. Suppose that $r_y \in b(A)$. Then $y \in A$, and, by assumption, A is a fixed set of ψ , so $A \in C(y)$; moreover, the elements $\{x_a : a \in A\}$ are products of reactions in $b(A)$. Thus, taking $U = \{x_a : a \in A\}$ we have $(U, r_y) \in \chi$; in other words, r_y is catalysed by products of reactions in $b(A)$. Since this holds for all $r_y \in b(A)$, it follows that $b(A)$ is a RAF.

Conversely, suppose that $b(W)$ is a RAF for \mathcal{Q}_ψ for some subset W of Y . Then for each $w \in W$, the reaction $r_w : f \rightarrow x_w$ has a catalyst set of the form $\{x_a : a \in A_w\}$ where A_w is a nonempty subset of W satisfying $w \in A_w$ and $\psi(A_w) = A_w$. It remains to show that $\psi(W) = W$. Since $\psi(W) \subseteq W$ it suffices to determine the converse containment. We have:

$$W \subseteq \bigcup_{w \in W} A_w = \bigcup_{w \in W} \psi(A_w) = \psi \left(\bigcup_{w \in W} A_w \right) \subseteq \psi(W),$$

as required to establish Claim 1.

Next observe that the RAFs for \mathcal{Q}_ψ are simply the nonempty fixed sets of φ (the maxRAF operator on R_Y for the CRS \mathcal{Q}_ψ), and so ψ and $b^{-1} \circ \varphi \circ b$ are two interior


operators on 2^Y , and these two functions the same fixed sets by Claim 1 (noting also that φ and ψ also trivially fix \emptyset). However, any two interior operators on 2^Y that have the same collection of fixed sets are identical functions. This is because any interior map ν on a set 2^Y is completely determined by its fixed sets since for any set $Y' \subseteq Y$ we have:

$$\nu(Y') = \bigcup \{U \subseteq Y' : \nu(U) = U\}.$$

Thus $\psi = b^{-1} \circ \varphi \circ b$, completing the proof. \square

Remark: It can be shown that Claim 1 also holds if we restrict the catalyst sets $U = \{x_a : a \in A\}$ to just those that correspond to the *minimal* sets $A \in C(y)$ (where minimal refers to set inclusion).


An example to illustrate Corollary 1: Consider the following CRS (based on Kauffman's binary polymer model with food set $F = \{0, 1, 00, 01, 10, 11\}$ from Fig. 1 of [41]):



This system is itself a RAF and it contains six other RAFs as subsets, namely $\{r_1, r_2\}, \{r_3\}, \{r_1, r_2, r_3\}, \{r_3, r_4\}, \{r_3, r_4, r_5\}, \{r_1, r_2, r_3, r_4\}$. Following the procedure described in the proof, we associate with r_i the reaction $r'_i : f[U_i] \rightarrow x_i$ where the set of (complex) catalysts is $U_i = \{\{x_j : r_j \in A\} : A \text{ is a RAF containing } r_i\}$. For example, for r_1 we have:

$$U_1 = \{\{x_1, x_2\}, \{x_1, x_2, x_3\}, \{x_1, x_2, x_3, x_4\}, \{x_1, x_2, x_3, x_4, x_5\}\}.$$

We can further simplify U_1 in line with remark following the proof of Proposition 1 above. For U_1 we only need to keep the single set $\{x_1, x_2\}$ since all other sets in U_1 contain this set. Applying this simplification to the other sets U_i leads to the resulting reduced system:



The RAFs of this system (with $F = \{f\}$) then correspond bijectively to the RAFs of the original system (under the bijection $r_i \leftrightarrow r'_i$ for each i).

Proof of Lemma 2: We establish the following more general result from which Lemma 2 follows by taking ψ to be the maxRAF operator (φ) for a CRS.

Given any function $\psi : 2^Y \rightarrow 2^Y$ define an associated derived function $\tilde{\psi} : 2^Y \rightarrow 2^Y$ as follows: For $X \subseteq Y$, let $F_\psi(X) = \{U \subseteq X : \psi(U) = U\}$ (the ‘fixed sets’ of ψ) and set:

$$\tilde{\psi}(X) = \bigcup \{U \in F_\psi(X) : \psi(U \setminus \{u\}) = \emptyset, \forall u \in U\}. \quad (5)$$

In words, $\tilde{\psi}(X)$ is the set of elements in X that lie in at least minimal fixed set of ψ .

Lemma 3. *If ψ is an interior operator on 2^Y then $\tilde{\psi}$ is also an interior operator on 2^Y , and $\tilde{\psi}(X) \subseteq \psi(X)$ for all $X \in 2^Y$.*

Proof: Condition (I_1) holds for $\tilde{\psi}$ since $\psi(X) \subseteq X$ for all $X \in 2^Y$ and so $U \in F_\psi(X)$ implies that $U \subseteq X$, and by definition $\tilde{\psi}(X)$ is a union of sets in $F_\psi(X)$. Condition (I_2) holds for $\tilde{\psi}$ since if $X \subseteq X'$ and $U \in F_\psi(X)$ then $U \in F_\psi(X')$. Thus it remains to establish (I_3). We have $\tilde{\psi}(\tilde{\psi}(X)) \subseteq \tilde{\psi}(X)$ by condition (I_1), and so it suffices to establish the reverse containment. Suppose that $U \in F_\psi(X)$. Then since $U \subseteq X$ and $\psi(U) = U$, it follows that $U = \psi(U) \subseteq \psi(X)$. Thus,

$$\{U \in F_\psi(X) : \psi(U \setminus \{u\}) = \emptyset, \forall u \in U\} \subseteq \{U \in F_\psi(\psi(X)) : \psi(U \setminus \{u\}) = \emptyset, \forall u \in U\},$$

and so the union of the sets on the left (i.e. $\tilde{\psi}(X)$) is a subset of the union of the sets on the right (i.e. $\tilde{\psi}(\tilde{\psi}(X))$). Thus, by Eqn. (5), (I_3) holds for $\tilde{\psi}$, and so $\tilde{\psi}$ is an interior operator on 2^Y .

The containment $\tilde{\psi}(X) \subseteq \psi(X)$ for all $X \in 2^Y$ follows by observing from Eqn. (5) that $\tilde{\psi}(X)$ is a union of a sub-collection of the set of subsets of X that are fixed by ψ , and the union of all such ψ -fixed subsets of X is $\psi(X)$. \square

Proof of Proposition 2: Suppose that $R_{\text{int}} := \bigcap_{i \geq 1} \hat{R}_i \neq \emptyset$. Then R_{int} is a RAF, and every reaction of R_{int} has a catalyst U_i that is not a subset of F , and so R_{int} is a strictly autocatalytic RAF. On the other hand, if R' is a strictly autocatalytic RAF, then since $R' \subseteq R$ we have (by induction on $i \geq 1$) $R' \subseteq \hat{R}_i$ for each i . Thus, since $R_{\text{int}} = \hat{R}_k$ for some value of k we have $R' \subseteq R_{\text{int}}$ and so R_{int} is the unique maximal strictly autocatalytic RAF for \mathcal{Q} . \square

Proof of Proposition 3: The argument just relies on the interior operator properties of φ established in Lemma 1. If $\varphi(\varphi(R) \setminus \{r_1, \dots, r_k\}) \neq \emptyset$ then $\varphi(R) \setminus \{r_1, \dots, r_k\}$ contains an iRAF, and this iRAF cannot be one of R_1, \dots, R_k , since for each i , r_i was chosen from R_i yet $r_i \notin \varphi(\varphi(R) \setminus \{r_1, \dots, r_k\})$ by definition. Conversely, if there is an iRAF R' of R that is different from R_1, \dots, R_k , then selecting r_i in $R_i \setminus R'$ for each i (this is possible since R_i is not a subset of R'), R' is an iRAF of $\varphi(R) \setminus \{r_1, \dots, r_k\}$ and so $R' = \varphi(R') \subseteq \varphi(\varphi(R) \setminus \{r_1, \dots, r_k\})$, thus this set is nonempty. \square

Proof of Proposition 4: First note that the problem described lies in the complexity class NP, since it can readily be verified (in polynomial time in $|\mathcal{Q}|$) whether or not a

given subset of reactions is an iRAF for \mathcal{Q} that contains a given reaction. We reduce an instance of the problem described to the following graph-theoretic problem: Given a finite directed graph $G = (V, A)$, and a vertex $v \in V$, determine whether or not there exist a chordless cycle in G that contains v . This problem was shown to be NP-complete in [31] (Theorem 1).

Given an arbitrary finite directed graph $G = (V, A)$, consider the following CRS $\mathcal{Q}_G = (X, R, \chi, F)$ where $F = \{f\}$, $X = V \cup F$, $R = \{r_v : v \in V\}$ where r_v is the reaction $r_v : f [u_1, \dots, u_n] \rightarrow v$ for each ordered pair $(u_i, v) \in A$. Notice that \mathcal{Q}_G is a CRS for which there is a single reactant for each reaction (namely f) which comprises the food set. Moreover, each reaction has singleton catalysts (i.e. complex catalysis is not involved).

Now any CRS $\mathcal{Q} = (X, R, \chi, F)$ that has the property that (i) each of its reactions has all its reactants in the food set, and (ii) each catalyst of each reaction is a singleton element in $X \setminus F$, then the iRAFs for \mathcal{Q} correspond precisely to the subsets of R that form a chordless cycle of the graph $G_{\mathcal{Q}} := (R, A)$ where $(r, r') \in A$ if a product of r is a catalyst of r' , (Theorem 2.1(ii) of [41]). It follows that the iRAFs for \mathcal{Q}_G are precisely the chordless cycles of G , which establishes the claimed reduction.

Proof of Proposition 5: Suppose that R' is a minimal subset of $\max\text{RAF}(\mathcal{Q})$ that is a RAF and produces all of the elements x_1, \dots, x_k . Then R' is a RAF of $\mathcal{Q}^+ := \mathcal{Q} + \{x_1, x_2, \dots, x_k\}$. Moreover, if R'' is a strict subset of R' that is both a RAF and that produces x_1, \dots, x_k , then R'' is also a RAF of \mathcal{Q}^+ and so R' is not a minimal set with the joint properties of being a RAF and producing x_1, \dots, x_k . Thus R' is an iRAF of \mathcal{Q}^+ . Conversely, suppose that R' is an iRAF of \mathcal{Q}^+ . Then R' is a RAF of \mathcal{Q} and R' must produce all of the elements x_1, \dots, x_k (since all reactions in \mathcal{Q}^+ require each of the elements x_1, \dots, x_k to be produced because none of these elements lie in the food set F (by assumption)). Moreover, R' is minimal subset of R that is a RAF for \mathcal{Q} and produces x_1, \dots, x_k ; for otherwise R' contains a strict subset with these properties and so would be a RAF of \mathcal{Q}^+ contradicting the assumption that R' is an iRAF of \mathcal{Q}^+ . \square

Proof of Proposition 6: We begin with a lemma.

Lemma 4. *Suppose that R' and $R'' \subsetneq R'$ are both RAFs for $\mathcal{Q} = (X, R, \chi, F)$. Then $R' \setminus R''$ is a RAF for $(X, R, \chi, F \cup \pi(R''))$.*

To see why this holds, simply observe that if o' is an admissible ordering for R' and o'' is the induced ordering of $R' \setminus R''$ then o'' is an admissible ordering of $R' \setminus R''$ for the CRS with expanded food set $(X, R, \chi, F \cup \pi(R''))$.

Returning to the proof of Proposition 6, observe that (ii) and (iii) equivalent (by the comments following the last lemma) so it suffices to show that (iii) implies (i), and, conversely, that (i) implies (iii). Suppose first that (iii) holds. Since R' equals the (disjoint) union of the sets \hat{R}_i we can select an admissible ordering of \hat{R}_i for each i , and extend this to an ordering of the reactions of R' in which all reactions from \hat{R}_i come before those of \hat{R}_{i+1} for each i . This ordering is an admissible ordering of R' , and each reaction in R' is either catalysed by an element of the food set or by the product of another reaction from R' . Thus R' is a RAF for \mathcal{Q} .

The proof of the converse direction ((i) implies (iii)) follows from Lemma 4, by induction on the length of the sequence R_1, R_2, \dots and the fact that every RAF contains an iRAF. \square